

The Open University's repository of research publications
and other research outputs

The Proteome and Citrullinome of Hippoglossus hippoglossus Extracellular Vesicles—Novel Insights into Roles of the Serum Secretome in Immune, Gene Regulatory and Metabolic Pathways

Journal Item

How to cite:

Magnadóttir, Bergljót; Kraev, Igor; Dodds, Alister W. and Lange, Sigrun (2021). The Proteome and Citrullinome of Hippoglossus hippoglossus Extracellular Vesicles—Novel Insights into Roles of the Serum Secretome in Immune, Gene Regulatory and Metabolic Pathways. International Journal of Molecular Sciences, 22(2), article no. 875.

For guidance on citations see [FAQs](#).

© [not recorded]



<https://creativecommons.org/licenses/by/4.0/>

Version: Version of Record

Link(s) to article on publisher's website:
<http://dx.doi.org/doi:10.3390/ijms22020875>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.



Article

The Proteome and Citrullinome of *Hippoglossus hippoglossus* Extracellular Vesicles—Novel Insights into Roles of the Serum Secretome in Immune, Gene Regulatory and Metabolic Pathways

Bergljót Magnadóttir ¹, Igor Kraev ², Alister W. Dodds ³ and Sigrun Lange ^{4,*}

¹ Institute for Experimental Pathology at Keldur, University of Iceland, Keldnavegur 3, 112 Reykjavik, Iceland; bergmagn@hi.is

² Electron Microscopy Suite, Faculty of Science, Technology, Engineering and Mathematics, Open University, Milton Keynes MK7 6AA, UK; igor.kraev@open.ac.uk

³ MRC Immunochemistry Unit, Department of Biochemistry, University of Oxford, Oxford OX1 3QU, UK; awdodds@gmail.com

⁴ Tissue Architecture and Regeneration Research Group, Department of Biomedical Sciences, University of Westminster, London W1W 6UW, UK

* Correspondence: S.Lange@westminster.ac.uk; Tel: +44-(0)207-911-5000

Citation: Magnadóttir, B.; Kraev, I.; Dodds, A.W.; Lange, S. The Proteome and Citrullinome of *Hippoglossus hippoglossus* Extracellular Vesicles—Novel Insights into Roles of the Serum Secretome in Immune, Gene Regulatory and Metabolic Pathways. *Int. J. Mol. Sci.* **2021**, *22*, 875. <https://doi.org/10.3390/ijms22020875>

Received: 24 December 2020

Accepted: 15 January 2021

Published: 16 January 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Abstract: Extracellular vesicles (EVs) are lipid bilayer vesicles which are released from cells and play multifaceted roles in cellular communication in health and disease. EVs can be isolated from various body fluids, including serum and plasma, and are usable biomarkers as they can inform health status. Studies on EVs are an emerging research field in teleost fish, with accumulating evidence for important functions in immunity and homeostasis, but remain to be characterised in most fish species, including halibut. Protein deimination is a post-translational modification caused by a conserved family of enzymes, named peptidylarginine deiminases (PADs), and results in changes in protein folding and function via conversion of arginine to citrulline in target proteins. Protein deimination has been recently described in halibut ontogeny and halibut serum. Neither EV profiles, nor total protein or deiminated protein EV cargos have yet been assessed in halibut and are reported in the current study. Halibut serum EVs showed a poly-dispersed population in the size range of 50–600 nm, with modal size of EVs falling at 138 nm, and morphology was further confirmed by transmission electron microscopy. The assessment of EV total protein cargo revealed 124 protein hits and 37 deiminated protein hits, whereof 15 hits were particularly identified in deiminated form only. Protein interaction network analysis showed that deimination hits are involved in a range of gene regulatory, immune, metabolic and developmental processes. The same was found for total EV protein cargo, although a far wider range of pathways was found than for deimination hits only. The expression of complement component C3 and C4, as well as pentraxin-like protein, which were identified by proteomic analysis, was further verified in EVs by western blotting. This showed that C3 is exported in EVs at higher levels than C4 and deiminated C3 was furthermore confirmed to be at high levels in the deimination-enriched EV fractions, while, in comparison, C4 showed very low detection in deimination-enriched EV fractions. Pentraxin was exported in EVs, but not detected in the deimination-enriched fractions. Our findings provide novel insights into EV-mediated communication in halibut serum, via transport of protein cargo, including post-translationally deiminated proteins.

Keywords: extracellular vesicles; proteome; citrullinome; peptidylarginine deiminase; deimination/citrullination; complement; pentraxin; immunity; metabolism; gene regulation

1. Introduction

Halibut is a teleost flatfish which belongs to the order Heterosomata (Pleuronectiformes). It is one of the largest teleost fish and endangered due to previous overfishing and slow rate of growth. The Atlantic halibut (*Hippoglossus hippoglossus* L.) is of considerable commercial value for aquaculture, where developmental abnormalities and viability in larval rearing have been one of the major obstacles [1,2]. Furthering understanding of immune, metabolic and developmental processes in commercially viable species, including halibut, is of great importance for the development of biomarkers associated to fish health and improved outcomes in aquaculture.

Peptidylarginine deiminases (PADs) are a calcium-dependent family of enzymes conserved throughout phylogeny with roles in physiological and pathophysiological processes [3–6]. PADs catalyse protein deimination/citrullination, which is an irreversible post-translational modification of protein arginine to citrulline, leading to structural and functional changes in target proteins [3,6,7]. Deimination can affect protein–protein interactions, as it modifies the protein structure and can cause protein denaturation or affect hydrogen bond formation [5,8]. Deimination can furthermore facilitate protein moonlighting, allowing one protein to carry out various functions within one polypeptide chain [9]. Intrinsically disordered proteins and β -sheets are most prone to undergo deimination and the position of the arginine within the protein plays roles as well [6,8,10]. While in fish, only one PAD form is present [11–14], mammals contain five tissue-specific PAD isozymes, with varying preferences for target proteins [3–5]. In other phyla, such as reptiles and birds, only three PAD forms are described [3,15,16], and PAD homologues are identified lower in the phylogeny tree [17], including in bacteria [18,19], fungi [20], parasites [21], as well as in Crustacea [22], Merostomata [23] and Mollusca [24]. PAD-mediated protein deimination has been reported in a range of taxa throughout the phylogeny tree, both in ontogeny, serum and plasma, as well as forming part of extracellular vesicle (EV) protein cargo [12–14,16,22–24].

EVs are lipid-bilayer vesicles in the size range of 50–1000 nm, released from most cells and participate in cellular communication in physiology and pathological processes. EVs are classified into small EVs (“exosomes”, <100 nm) and larger EVs (“microvesicles” 100–1000 nm), which are released from cells via different biogenesis pathways, including exocytosis or membrane blebbing [25,26]. Roles for PADs in the modulation of EV release have furthermore been described [27–29]. EVs carry a range of cargo, including proteins, enzymes, genetic material, long non-coding RNAs and microRNAs, derived from the cells of origin [25–33]. Protein EV cargo can furthermore consist of post-translationally modified proteins, which possibly contribute differently to cellular communication compared with non-modified protein forms. Therefore, it may be of considerable interest to gain insight into differences in such protein cargo in serum-EVs to further understanding of post-translational modifications (PTMs) in cellular communication.

While EV research has been an exponentially expanding field in the past decade in relation to human disease, less is known about EV communication in other taxa. The comparative field of EV research has recently been growing, including by studies from our group [14,16,19,22–24,32–40]. Therefore, there is currently great interest in expanding EV studies, also in relation to teleost fish and biomarker discovery for aquaculture [32,33,38,41]. Furthermore, fundamental research into EV communication across the phylogeny tree will allow for increased understanding of EV-mediated pathways in evolution.

This study aimed at characterising EVs from halibut sera, assessing both total proteomic cargo and deiminated protein cargo to gain insights into putative roles for protein deimination in the serum secretome.

2. Results

2.1. EV Profiling from Halibut Sera

Halibut serum EVs were characterised by NTA, revealing a poly-dispersed EV population in the size range of 50–600 nm, with the modal size of EVs falling at 138 nm (Figure 1A). The EVs were further characterised for two EV specific markers, CD63 and Flotillin-1 and found positive for both (Figure 1B). EV morphology was further confirmed by transmission electron microscopy (TEM), revealing typical EV morphology (see arrows) and confirming a polydispersed population (Figure 1C).




Figure 1. Halibut serum-extracellular vesicles (EV)s were characterised by: (A) Nanoparticle tracking analysis (NTA), showing size distribution profiles of EVs in the size range of 50–600 nm, with the modal size of vesicles at 138 nm; (B) Western blotting (WB) analysis shows that the EVs are positive for CD63 and Flotillin-1; (C) Transmission electron microscopy (TEM) showing EV morphology—see arrows pointing at EVs (scale bar is indicated at 20 nm).

2.2. The Proteome and Citrullinome of Halibut Serum EVs

Total protein content, as well as F95 enriched protein content, representative of deiminated protein cargo in EVs (the “EV-citrullinome”), was identified by LC-MS/MS analysis. A range of proteins relating to innate and adaptive immunity, as well as gene regulation and cellular function, were identified as deiminated in EV cargo, and are listed in Table 1 (for full details on LC-MS/MS analysis, see Supplementary Table S1). Total EV protein cargo analysis revealed proteins relating to innate and adaptive immunity, nuclear proteins relating to gene regulation, proteins relating to cellular function and metabolism and are listed in Table 2 (for full details on LC-MS/MS analysis, see Supplementary Table S2). Total serum-EV proteins stained by silver staining are shown in Figure 2A, F95 enriched proteins from serum-EVs are shown in Figure 2B and the number of total EV proteins identified, overlapping with deiminated/citrullinated EV proteins identified are presented in the Venn diagram in Figure 2C.

Table 1. Deiminated proteins in serum extracellular vesicles (EVs) of halibut (*Hippoglossus hippoglossus* L), as identified by F95-enrichment in conjunction with LC-MS/MS analysis. Deiminated proteins were isolated from serum-EVs from a pool of $n = 4$ fish, using immunoprecipitation with the pan-deimination F95 antibody. The resulting F95-enriched eluate was then analysed by LC-MS/MS and peak list files submitted to Mascot, using the Teleost UniProt database. Peptide sequence hits are listed, showing the number of sequences for protein hits and total score. Species hit names are indicated. In the case of uncharacterised protein ID, proteins matching the same set of peptides are indicated in brackets. Protein hits highlighted in pink (*) are specific to the F95 enriched EV fraction only. A full list of protein sequence hits and peptides is further provided in Supplementary Table S1.

Protein ID Protein Name	Species Name Common Name	Matches (Sequences)	Total Score ($p < 0.05$) [†]
A0A6J2W3P0_CHACN Uncharacterised protein (histone H3-like)	<i>Chanos chanos</i> Milkfish	16 (13)	538
A0A672ZYE0_9TELE Uncharacterised protein	<i>Sphaeramia orbicularis</i> Orbiculate cardinalfish	12 (9)	451
A0A0A1G3Q1_9TELE Beta-actin	<i>Oxyeleotris marmorata</i> Marble goby	10 (10)	427
A0A3P8Y5X6_ESOLU IF rod domain-containing protein	<i>Esox Lucius</i> Northern pike	26 (8)	356
*W5ZLY1_9TELE Cytoplasmic 2 actin	<i>Campylomormyrus compressirostris</i> Elephantfish	8 (8)	336
A0A3B4ZTX8_9TELE Uncharacterized protein (NTR domain-containing protein; Complement component C3)	<i>Stegastes partitus</i> Bicolour damselfish	8 (7)	324
A0A3B4THR8_SERDU Uncharacterized protein (NTR domain-containing protein; anaphylatoxin-like, Complement component C3)	<i>Seriola dumerili</i> Greater amberjack	9 (8)	299
A0A6G0HQ07_LARCR Histone H4	<i>Larimichthys crocea</i> Yellow croaker	8 (6)	281
A0A3Q3IVX9_MONAL Uncharacterized protein (Complement C3)	<i>Monopterus albus</i> Asian swamp eel	8 (7)	276
A0A3P9BEG5_9CICH Uncharacterized protein (Anaphylatoxin-like, complement C3)	<i>Maylandia zebra</i> Zebra mbuna	6 (6)	273
A0A484CCU5_PERFV Uncharacterized protein (complement C3)	<i>Perca flavescens</i> Yellow perch	8 (7)	271
A5JV31_HIPHI Phosvitin	<i>Hippoglossus hippoglossus</i> Atlantic halibut	7 (7)	261
*A0A087XQB5_POEFO Tubulin alpha chain	<i>Poecilia formosa</i> Amazon molly	6 (5)	256
A0A6F9CZC7_9TELE Uncharacterized protein (tubulin alpha-chain)	<i>Coregonus</i> sp. 'balchen' Whitefish, salmonidae	5 (4)	251
*Q1RLR3_DANRE Keratin 93	<i>Danio rerio</i> Zebrafish	8 (5)	237
A0A1S5XZE7_9TELE Histone H3	<i>Lipogramma levinsoni</i> Hourglass basslet	7 (7)	231
A3F5V1_ORENI Beta actin (Fragment)	<i>Oreochromis niloticus</i> Nile tilapia	7 (7)	222
A0A5N5KJN7_PANHP IF rod domain-containing protein	<i>Pangasianodon hypophthalmus</i> Iridescent shark	6 (4)	185
A0A4W6CP97_LATCA Uncharacterized protein (Alpha-2-macroglobulin)	<i>Lates calcarifer</i> Barramundi/Asian sea bass	5 (4)	179
*H2MSJ5_ORYLA	<i>Oryzias latipes</i>	5 (4)	159

Uncharacterized protein	Medaka/Japanese rice fish		
A0A060WDP8_ONCMY	<i>Oncorhynchus mykiss</i>	3 (3)	136
Elongation factor 1- α	Rainbow trout		
A0A671UYU7_SPAAU	<i>Sparus aurata</i>		
Uncharacterized protein (A2M_recep domain-containing protein)	Gilt-head bream	3 (1)	117
G3Q4A0_GASAC	<i>Gasterosteus aculeatus</i>	2 (2)	116
Fibrinogen beta chain	Three-spined stickleback		
A0A0F8AH88_LARCR	<i>Larimichthys crocea</i>	2 (2)	107
Ig heavy chain V region 5A	Yellow croaker		
A0A4W6FLR7_LATCA	<i>Lates calcarifer</i>		
Uncharacterized protein (NTR domain-containing protein; anaphylatoxin like; A2M_N_2 domain-containing; complement C5)	Barramundi/Asian sea bass	3 (3)	104
A0A4Z2B138_9TELE	<i>Takifugu bimaculatus</i>	3 (3)	99
Anaphylatoxin-like domain-containing protein	Pufferfish		
Q4KVK3_HIPHI	<i>Hippoglossus hippoglossus</i>	2 (2)	94
Complement component c3 (Fragment)	Atlantic halibut		
*A0A5J5C7F1_9PERO	<i>Etheostoma spectabile</i>	2 (2)	94
Uncharacterized protein (Fragment)	Orangethroat darter		
*A0A0P7WL38_SCLFO	<i>Scleropages formosus</i>	4 (2)	93
Trypsin-3-like	Asian arowana		
Q5DVG8_PLAFE	<i>Platichthys flesus</i>	3 (2)	84
Apolipoprotein AI	European flounder		
A0A0F8ABH4_LARCR	<i>Larimichthys crocea</i>	3 (1)	82
Granzyme B(G,H)	Yellow croaker		
A0A484D989_PERFV	<i>Perca flavescens</i>	3 (2)	71
Peptidase S1 domain-containing protein	Yellow perch		
*A0A5N5Q536_PANHP	<i>Pangasianodon hypophthalmus</i>	2 (2)	70
Centrosomal protein of 162 kDa	Iridescent shark		
*A0A0P7UEW6_SCLFO	<i>Scleropages formosus</i>	1 (1)	69
2-phospho-D-glycerate hydro-lyase	Asian arowana		
A0A060YWU0_ONCMY	<i>Oncorhynchus mykiss</i>	4 (2)	68
Peptidase S1 domain-containing protein	Rainbow trout		
*A0A1A7WRH6_9TELE	<i>Iconisemion striatum</i>	2 (2)	64
Integrin beta	Killifish		
A0A3B5M528_9TELE	<i>Xiphophorus couchianus</i>	1 (1)	64
Serotransferrin	Monterrey platyfish		
A0A060Z3N3_ONCMY	<i>Oncorhynchus mykiss</i>	2 (2)	63
Ig-like domain-containing protein	Rainbow trout		
A0A060W543_ONCMY	<i>Oncorhynchus mykiss</i>	2 (2)	62
Histone H2A	Rainbow trout		
A0A0R4IU44_DANRE	<i>Danio rerio</i>	1 (1)	61
Inter-alpha-trypsin inhibitor heavy chain 3b	Zebrafish		
HV05_CARAU	<i>Carassius auratus</i>	2 (1)	60
Ig heavy chain V region 5A	Goldfish		
*A0A060XD44_ONCMY	<i>Oncorhynchus mykiss</i>	4 (2)	60
Uncharacterized protein	Rainbow trout		
A0A4W5L5T6_9TELE	<i>Hucho hucho</i>	1 (1)	57
Thioredoxin	Danube salmon		
*A0A3Q3LZB0_9TELE	<i>Mastacembelus armatus</i>	1 (1)	57
Uncharacterized protein	Zig-zag eel/Spiny eel		
*A0A5J5DS23_9PERO	<i>Etheostoma spectabile</i>	1 (1)	57
Uncharacterized protein	Orangethroat darter		
*A0A3B3QST7_9TELE	<i>Paramormyrops kingsleyae</i>	1 (1)	55
Uncharacterized protein	Elephantfish		
*A0A0E9RVI6_ANGAN	<i>Anguilla Anguilla</i>	1 (1)	53

Uncharacterized protein	European eel		
*A0A3Q3SSB4_9TELE	<i>Mastacembelus armatus</i>	1 (1)	53
*Myosin_tail_1 domain-containing protein	Zig-zag eel/Spiny eel		

[†] Ions score is $-10 \times \log(P)$, where P is the probability that the observed match is a random event. Individual ions scores > 53 indicate identity or extensive homology ($p < 0.05$). Protein scores are derived from ions scores as a non-probabilistic basis for ranking protein hits.

Table 2. Total protein cargo in serum-EVs of halibut (*Hippoglossus hippoglossus* L), as identified by LC-MS/MS analysis from serum-EVs isolated from a pool of sera from $n = 4$ fish. Peak list files were submitted to Mascot, using the Teleost UniProt database. Peptide sequence hits are listed, showing the number of sequences for protein hits and total score. Species hit names are indicated. In the case of uncharacterised protein ID, proteins matching the same set of peptides are indicated in brackets. Protein hits highlighted in blue (*) were not identified in the F95 enriched fraction. A full list of protein sequence hits and peptides is further provided in Supplementary Table S2.

Protein ID Protein Name	Species Name Common Name	Matches (Sequences)	Total Score ($p < 0.05$) [†]
A5JV31_HIPHI <i>Phosvitin</i>	<i>Hippoglossus hippoglossus</i> Atlantic halibut	145 (56)	3616
A5JV30_HIPHI <i>Phosvitin</i>	<i>Hippoglossus hippoglossus</i> Atlantic halibut	90 (52)	3303
Q4KVK3_HIPHI Complement component c3 (fragment)	<i>Hippoglossus hippoglossus</i> Atlantic halibut	69 (25)	1690
A0A2U9BPE5_SCOMX Complement component C3 isoform 2	<i>Scophthalmus maximus</i> Turbot	89 (24)	1426
A0A3B4THR8_SERDU Uncharacterized protein (NTR domain-containing protein, Complement C3-like, A2M_recep domain-containing protein)	<i>Seriola dumerili</i> Greater amberjack	79 (22)	1269
A0A3B4TYC3_SERDU NTR domain-containing protein	<i>Seriola dumerili</i> Greater amberjack	65 (21)	1250
Q9PTY1_PAROL Complement component C3	<i>Paralichthys olivaceus</i> Olive flounder	70 (22)	1176
G4WAB7_EPICO Complement component c3	<i>Epinephelus coioides</i> Orange-spotted grouper	57 (20)	1145
A0A3P9BEG5_9CICH Uncharacterized protein (Anaphylatoxin-like domain-containing protein; C3a)	<i>Maylandia zebra</i> Zebra mbuna	66 (19)	1120
*A0A669BPJ4_ORENI Uncharacterized protein	<i>Oreochromis niloticus</i> Nile tilapia	71 (18)	1097
A0A671YHA0_SPAAU Uncharacterized protein (C3)	<i>Sparus aurata</i> Gilt-head bream	45 (17)	904
*A0A6A5FQW4_PERFL Uncharacterized protein	<i>Perca fluviatilis</i> European perch	57 (16)	885
A0A484CCU5_PERFV Uncharacterized protein (Anaphylatoxin-like domain-containing protein)	<i>Perca flavescens</i> Yellow perch	56 (17)	879
F8R8R1_DICLA Complement component c3-2	<i>Dicentrarchus labrax</i> European bass	59 (15)	871
A0A484DL37_PERFV Anaphylatoxin-like domain-containing protein	<i>Perca flavescens</i> Yellow perch	42 (15)	784
A0A4W6E087_LATCA Complement component c3a, duplicate 5	<i>Lates calcarifer</i> Barramundi/Asian sea bass	43 (15)	744
A0A6A5FJW4_PERFL Uncharacterized protein (Integrase catalytic domain-containing protein, Alpha-2-macroglobulin-like)	<i>Perca fluviatilis</i> European perch	14 (12)	600
A0A2P9DTV2_SOLSE <i>Phosvitin</i>	<i>Solea senegalensis</i> Senegalese sole	16 (10)	594
Q6QZI2_PSEAM	<i>Pseudopleuronectes americanus</i>	37 (9)	574

Complement component C3 (Fragment)	Winter flounder		
A0A3Q1ID66_ANATE	<i>Anabas testudineus</i>	25 (9)	569
Phosvitin	Climbing perch		
*A0A4W6F6V9_LATCA	<i>Lates calcarifer</i>	15 (9)	549
Apolipoprotein Bb, tandem duplicate 2	Barramundi/Asian sea bass		
A0A6G1PAV1_9TELE	<i>Channa argus</i>		
Complement C3 Complement C3 beta chain Complement C3 alpha chain	Northern snakehead	38 (10)	540
*A0A4P8JD10_9TELE	<i>Lateolabrax maculatus</i>	13 (9)	532
Apolipoprotein Bb.1	Spotted sea bass		
*A0A6A5DT05_PERFL	<i>Perca fluviatilis</i>	16 (9)	529
Vitellogenin domain-containing protein	European perch		
A0A673IJP2_9TELE	<i>Sinocyclocheilus rhinoceros</i>		
IF rod domain-containing protein	<i>Sinocyclocheilus cavefish</i> (Cypri- nid)	48 (12)	528
A0A4W6CMC4_LATCA	<i>Lates calcarifer</i>		
Uncharacterized protein (Alpha-2-macroglobulin)	Barramundi/Asian sea bass	14 (11)	525
A0A3P8Y5X6_ESOLU	<i>Esox Lucius</i>		
IF rod domain-containing protein	Northern pike	51 (11)	499
A0A6G1PQL3_9TELE	<i>Channa argus</i>		
Alpha-2-macroglobulin	Northern snakehead	12 (10)	497
A0A6A4SX26_SCOMX	<i>Scophthalmus maximus</i>		
IF rod domain-containing protein	Turbot	51 (11)	463
Q5DVG8_PLAFE	<i>Platichthys flesus</i>		
Apolipoprotein AI	European flounder	26 (9)	453
*A0A3B4T6U1_SERDU	<i>Seriola dumerili</i>	12 (9)	440
Vitellogenin domain-containing protein	Greater amberjack		
A0A665VQL3_ECHNA	<i>Echeneis naucrates</i>		
Uncharacterized protein (A2M_N_2 domain-containing protein)	Live sharksucker	9 (8)	409
*A0A2U9D044_SCOMX	<i>Scophthalmus maximus</i>		
Putative apolipoprotein B-100-like isoform 2	Turbot	14 (8)	407
*Q9PVW6_PAROL	<i>Paralichthys olivaceus</i>		
Complement component C9	Olive flounder	14 (7)	403
A0A4W6FLR7_LATCA	<i>Lates calcarifer</i>		
Uncharacterized protein (Anaphylatoxin-like domain-containing, A2M_N_2 domain containing protein, NTR domain containing protein, Complement C5)	Barramundi/Asian sea bass	10 (8)	386
A0A4W6CP97_LATCA	<i>Lates calcarifer</i>		
Uncharacterized protein (A2M_recep domain-containing protein, TED_complement domain-containing protein)	Barramundi/Asian sea bass	17 (7)	362
*A0A3P8RR96_AMPPE	<i>Amphiprion percula</i>		
Complement component C9	Orange clownfish	12 (5)	353
A0A3Q1HZ43_ANATE	<i>Anabas testudineus</i>		
Uncharacterized protein (Inter-alpha-trypsin inhibitor, VIT domain-containing protein)	Climbing perch	13 (9)	336
*A0A3Q1H6Y9_ANATE	<i>Anabas testudineus</i>		
Complement component 8 subunit beta	Climbing perch	8 (8)	336
A0A6G1PI27_9TELE	<i>Channa argus</i>		
Inter-alpha-trypsin inhibitor heavy chain H3	Northern snakehead	13 (7)	324
A0A6A5FLM2_PERFL	<i>Perca fluviatilis</i>		
Uncharacterized protein (alpha-2-macroglobulin-like, A2M_recep domain-containing protein)	European perch	10 (6)	323
A0A6A5FFR2_PERFL	<i>Perca fluviatilis</i>		
Anaphylatoxin-like domain-containing protein	European perch	15 (7)	323
A0A484DIJ5_PERFV	<i>Perca flavescens</i>		
Uncharacterized protein (Alpha-2-macroglobulin)	Yellow perch	11 (7)	321

A0A6A5FE70_PERFL	<i>Perca fluviatilis</i>		
Uncharacterized protein (A2M_recep domain-containing, MG2 domain-containing protein)	European perch	11 (7)	318
A0A6J2W3P0_CHACN	<i>Chanos chanos</i>		
uncharacterized protein LOC115819396 (Histone H4, Histone H3, Histone H2B)	Milkfish	8 (7)	312
*A0A665V532_ECHNA	<i>Echeneis naucrates</i>	8 (6)	310
Plasminogen	Live sharksucker		
A0A3Q3L7G2_9TELE	<i>Mastacembelus armatus</i>	6 (6)	308
Complement component c3b, tandem duplicate 2	Zig-zag eel/Spiny eel		
*CO8B_PAROL	<i>Paralichthys olivaceus</i>	5 (5)	304
Complement component C8 beta chain	Olive flounder		
A0A671PIL3_9TELE	<i>Sinocyclocheilus anshuiensis</i>	17 (6)	301
IF rod domain-containing protein	Sinocyclocheilus cavefish (Cyprinoid)		
*A0A3Q3E5X5_9LABR	<i>Labrus bergylta</i>	7 (4)	298
Uncharacterized protein (C1q domain-containing protein)	Ballan wrasse		
*A0A3Q0S0V4_AMPPI	<i>Amphilophus citrinellus</i>	18 (5)	292
Uncharacterized protein	Midas cichlid		
A0A6A4SU52_SCOMX	<i>Scophthalmus maximus</i>	7 (7)	291
Uncharacterized protein (Complement component c3b)	Turbot		
*A0A3P8TA20_AMPPE	<i>Amphiprion percula</i>	11 (7)	290
Zgc:112265	Orange clownfish		
A0A096MDQ7_POEFO	<i>Poecilia formosa</i>	11 (6)	288
Phosvitin	Amazon molly		
Q5XVQ2_FUNHE	<i>Fundulus heteroclitus</i>	17 (5)	288
Apolipoprotein A1 (Fragment)	Atlantic killifish, mud minnow		
*Q6QZ19_PSEAM	<i>Pseudopleuronectes americanus</i>	12 (5)	284
Complement component C9 (Fragment)	Winter flounder		
*A0A4U5UPP9_COLLU	<i>Collichthys lucidus</i>	7 (5)	283
Apolipoprotein B-100	(Big head croaker)		
A0A3Q1EMN2_9TELE	<i>Acanthochromis polyacanthus</i>	8 (5)	280
Uncharacterized protein (beta actin, actin cytoplasmic-1)	Spiny chromis damselfish		
*A0A6J2P874_COTGO	<i>Cottoperca gobio</i>	7 (5)	280
plasminogen	Channel bull blenny		
A0A3B4VID4_SERDU	<i>Seriola dumerili</i>	9 (5)	279
Uncharacterized protein (MG2 domain-containing protein)	Greater amberjack		
*A0A3Q3M9S2_9TELE	<i>Mastacembelus armatus</i>	12 (4)	278
Uncharacterized protein	Zig-zag eel/Spiny eel		
W5ZMG9_9TELE	<i>Campylomormyrus compressirostris</i>	7 (4)	267
Cytoplasmic 1 actin	Elephantfish		
A0A553Q7M4_9TELE	<i>Danionella translucida</i>	6 (6)	262
Uncharacterized protein (Histone H2A, H2B putative, H3)	Micro glassfish (Cyprinid)		
A0A3Q1H0X2_ANATE	<i>Anabas testudineus</i>	5 (5)	260
Complement component c3b, tandem duplicate 2	Climbing perch		
*A0A6A4SHP5_SCOMX	<i>Scophthalmus maximus</i>	12 (5)	258
Uncharacterized protein	Turbot		
*G3NNM8_GASAC	<i>Gasterosteus aculeatus</i>	6 (6)	256
Uncharacterized protein	Three-spined stickleback		
*A0A0P7YVM9_SCLFO	<i>Scleropages formosus</i>	10 (5)	251
Keratin, type I cytoskeletal 13-like	Asian arowana		
*A0A6A4SWR2_SCOMX	<i>Scophthalmus maximus</i>	7 (6)	251
EGF-like domain-containing protein	Turbot		
A0A2U9B3I5_SCOMX	<i>Scophthalmus maximus</i>	13 (6)	247
Alpha-2-macroglobulin	Turbot		
A0A4Z2BCD9_9TELE	<i>Takifugu bimaculatus</i>	6 (5)	242
Uncharacterized protein	Pufferfish		

(Complement C5 C3 and PZP-like alpha-2-macroglobulin domain-containing protein)

A0A671TD78_SPAAU	<i>Sparus aurata</i>		
Complement component c3b, tandem duplicate 2	Gilt-head bream	5 (5)	238
*A0A0A0QKL5_OPLFA	<i>Oplegnathus fasciatus</i>		
Complement component 4	Striped beakfish	6 (5)	234
*A0A6A4RUD7_SCOMX	<i>Scophthalmus maximus</i>		
Vitellogenin domain-containing protein	Turbot	6 (6)	233
A0A672YA60_9TELE	<i>Sphaeramia orbicularis</i>		
Uncharacterized protein (inter-alpha-trypsin inhibitor heavy chain)	Orbiculate cardinalfish	7 (6)	232
*A0A672JL95_SALFA	<i>Salarias fasciatus</i>		
Uncharacterized protein (complement C7)	Lawnmower blenny	5 (5)	232
*A0A3B4Y8X6_SERLL	<i>Seriola lalandi</i>		
Uncharacterized protein (Hephaestin-like protein 1, Desmoglein-2)	Yellowtail amberjack	10 (4)	231
*A0A3B4UHS2_SERDU	<i>Seriola dumerili</i>		
Uncharacterized protein	Greater amberjack	4 (4)	229
*A0A087YMZ0_POEFO	<i>Poecilia formosa</i>		
Uncharacterized protein (Ceruloplasmin)	Amazon molly	11 (6)	229
A0A3Q4G4S3_NEOBR	<i>Neolamprologus brichardi</i>		
Uncharacterized protein (NTR domain-containing protein)	Lyretail cichlid	11 (5)	229
*A0A3Q1EBE7_9TELE	<i>Acanthochromis polyacanthus</i>		
Vitellogenin domain-containing protein	Spiny chromis damselfish	4 (4)	228
A0A3P9A8D3_ESOLU	<i>Esox Lucius</i>		
Uncharacterized protein (Alpha-2-macroglobulin, A2M_recep domain-containing)	Northern pike	8 (4)	218
*A0A3P8WZ01_CYNSE	<i>Cynoglossus semilaevis</i>		
Vitellogenin domain-containing protein	Tongue sole	7 (4)	217
*A0A3B4F9T0_9CICH	<i>Pundamilia nyererei</i>		
Carboxypeptidase Q	Cichlid	3 (3)	215
*A0A6J2QSS9_COTGO	<i>Cottoperca gobio</i>		
complement component C9	Channel bull blenny	3 (2)	209
*A0A672GNQ4_SALFA	<i>Salarias fasciatus</i>		
Vitellogenin domain-containing protein	Lawnmower blenny	7 (4)	208
*A0A3B4ULR2_SERDU	<i>Seriola dumerili</i>		
Zgc:112265	Greater amberjack	9 (5)	207
A0A3B4THN2_SERDU	<i>Seriola dumerili</i>		
Fibrinogen beta chain	Greater amberjack	4 (4)	205
*A0A2U9BK85_SCOMX	<i>Scophthalmus maximus</i>		
Putative complement component C8 alpha chain	Turbot	3 (3)	203
*G8DP14_PLAFE	<i>Platichthys flesus</i>		
Beta 1-globin	European flounder	4 (4)	201
*A0A0F8C5A6_LARCR	<i>Larimichthys crocea</i>		
Antithrombin-III	Yellow croaker	6 (5)	200
*A0A2U9CEJ2_SCOMX	<i>Scophthalmus maximus</i>		
Complement component 7	Turbot	4 (4)	200
A0A5C6MX12_9TELE	<i>Takifugu flavidus</i>		
Complement C3	Yellowbelly pufferfish	17 (5)	196
*Q6QZI5_PSEAM	<i>Pseudopleuronectes americanus</i>		
Complement component C8 beta chain	Winter flounder	4 (4)	194
*A0A3B3BJ38_ORYME	<i>Oryzias melastigma</i>		
Vitellogenin domain-containing protein	Marine medaka	7 (4)	192
*A0A6J2S534_COTGO	<i>Cottoperca gobio</i>		
apolipoprotein B-100	Channel bull blenny	5 (5)	190
*A0A3Q1JFY5_ANATE	<i>Anabas testudineus</i>	5 (3)	187

Uncharacterized protein (ceruloplasmin)	Climbing perch		
A0A672I1M9_SALFA	<i>Salarias fasciatus</i>		
Uncharacterized protein (Inter-alpha-trypsin inhibitor heavy chain, VIT domain-containing protein)	Lawnmower blenny	6 (4)	186
A0A3B5AT07_9TELE	<i>Stegastes partitus</i>	7 (4)	185
IF rod domain-containing protein	Bicolour damselfish		
*A0A4Z2CEC7_9TELE	<i>Takifugu bimaculatus</i>	4 (4)	183
Uncharacterized protein (complement C4)	Pufferfish		
*A0A3Q3II57_MONAL	<i>Monopterus albus</i>	5 (4)	183
Uncharacterized protein	Asian swamp eel		
*A0A3Q3FIH8_KRYMA	<i>Kryptolebias marmoratus</i>	7 (4)	180
Uncharacterized protein	Mangrove rivulus(killifish)		
*A0A2U9AYP3_SCOMX	<i>Scophthalmus maximus</i>	5 (3)	177
Complement component 4	Turbot		
*A0A6J2RDF1_COTGO	<i>Cottoperca gobio</i>	4 (4)	176
complement C4-B-like	Channel bull blenny		
A0A4W6ERJ2_LATCA	<i>Lates calcarifer</i>	5 (4)	173
Fibrinogen gamma chain	Barramundi/Asian sea bass		
*A0A2I4C034_9TELE	<i>Austrofundulus limnaeus</i>	3 (3)	167
collagen alpha-1(XII) chain	Killifish		
*A0A6I9PPD4_9TELE	<i>Notothenia coriiceps</i>	4 (4)	167
complement C4-like	Black rockcod/Antarctic yel-lowbelly rockcod		
*H3BWT7_TETNG	<i>Tetraodon nigroviridis</i>	5 (3)	163
Ceruloplasmin	Green spotted puffer		
*Q4SXM5_TETNG	<i>Tetraodon nigroviridis</i>	5 (4)	160
Chromosome 12 SCAF12357, whole genome shotgun sequence	Green spotted puffer		
A0A1A8F2V0_9TELE	<i>Nothobranchius korthausae</i>	5 (2)	160
Uncharacterized protein (Alpha2-macroglobulin)	Killifish		
*A0A3B5BD88_9TELE	<i>Stegastes partitus</i>	4 (4)	159
Vitellogenin domain-containing protein	Bicolour damselfish		
A0A6G1QB31_9TELE	<i>Channa argus</i>	9 (2)	159
Serotransferrin	Northern snakehead		
*A0A060WU48_ONCMY	<i>Oncorhynchus mykiss</i>	2 (2)	157
Uncharacterized protein (Desmoplakin)	Rainbow trout		
*A0A3Q3FAE5_9LABR	<i>Labrus bergylta</i>	4 (3)	155
Complement component 8 subunit beta	Ballan wrasse		
A0A6J2Q526_COTGO	<i>Cottoperca gobio</i>	4 (3)	155
fibrinogen gamma chain	Channel bull blenny		
*A0A3B4UV22_SERDU	<i>Seriola dumerili</i>	6 (4)	154
Antithrombin-III	Greater amberjack		
*A0A3Q2QAA5_FUNHE	<i>Fundulus heteroclitus</i>	4 (3)	154
Uncharacterized protein	Atlantic killifish, mud minnow		
*A0A6J2P7B9_COTGO	<i>Cottoperca gobio</i>	3 (2)	153
apolipoprotein B-100-like	Channel bull blenny		
*A0A484D0P7_PERFV	<i>Perca flavescens</i>	6 (5)	153
Uncharacterized protein (ceruloplasmin)	Yellow perch		
*A0A3B4TA89_SERDU	<i>Seriola dumerili</i>	3 (3)	149
Uncharacterized protein	Greater amberjack		
*A0A673XMC1_SALTR	<i>Salmo trutta</i>	3 (3)	148
Uncharacterized protein (complement C4, C4-B)	Brown trout		
F8U8N8_CHELB	<i>Chelon labrosus</i>	4 (3)	146
Alpha 2 macroglobulin (fragment)	Thicklip grey mullet		
F2Y9S5_MORSA	<i>Morone saxatilis</i>	3 (3)	145
Phosvitin	Striped bass		
*A0A3P9Q7U6_POERE	<i>Poecilia reticulata</i>	4 (4)	144
Complement component C9	Guppy		

A0A0F8AH88_LARCR	<i>Larimichthys crocea</i>	9 (3)	143
Ig heavy chain V region 5A	Yellow croaker		
*A0A667Y3E0_9TELE	<i>Myripristis murdjan</i>	6 (3)	142
Vitellogenin domain-containing protein	Blacktipped soldierfish		
*A0A672QEF7_SINGR	<i>Sinocyclocheilus graham</i>	8 (4)	141
Uncharacterized protein	Golden-line barbell		
*A0A3B5B7I8_9TELE	<i>Stegastes partitus</i>	5 (4)	141
Antithrombin-III	Bicolour damselfish		
*A0A0B6VKQ1_ORYCL	<i>Oryzias celebensis</i>	3 (3)	139
B5 protein	Celebes medaka		
*A0A671TKG8_SPAAU	<i>Sparus aurata</i>	4 (2)	138
Uncharacterized protein	Gilt-head bream		
*A0A4P8JCG0_9TELE	<i>Lateolabrax maculatus</i>	3 (2)	136
Apolipoprotein Bb.2	Spotted sea bass		
*A0A3B4FS46_9CICH	<i>Pundamilia nyererei</i>	4 (3)	132
IGv domain-containing protein	Cichlid		
A0A3P9H4Z3_ORYLA	<i>Oryzias latipes</i>		
Uncharacterized protein (A2M_N_2 domain-containing protein, anaphylatoxin-like domain)	Medaka/Japanese rice fish	9 (3)	132
A0A0F8AKQ4_LARCR	<i>Larimichthys crocea</i>	5 (3)	131
Alpha-2-macroglobulin	Yellow croaker		
A0A3B4TIN1_SERDU	<i>Seriola dumerili</i>	3 (3)	130
Phosvitin	Greater amberjack		
B6RUP0_ORYDN	<i>Oryzias dancena</i>	4 (3)	129
Beta-actin (Fragment)	Indian ricefish		
*A0A484CD54_PERFV	<i>Perca flavescens</i>	3 (3)	129
Uncharacterized protein (Complement C7)	Yellow perch		
A0A3Q4FXR7_NEOBR	<i>Neolamprologus brichardi</i>	4 (3)	128
Ig-like domain-containing protein	Lyretail cichlid		
Q5SET8_9TELE	<i>Bembras japonica</i>	3 (3)	128
Histone H3 (Fragment)	Red flathead		
A0A3Q1IXI9_ANATE	<i>Anabas testudineus</i>	3 (3)	128
Uncharacterized protein (A2M_recep domain-containing protein)	Climbing perch		
*A0A4Z2H8W0_9TELE	<i>Liparis tanakae</i>	2 (2)	126
Biotinidase	Tanaka's snailfish		
A0A6G1PSN0_9TELE	<i>Channa argus</i>	6 (5)	126
Alpha-2-macroglobulin	Northern snakehead		
A0A669DKF1_ORENI	<i>Oreochromis niloticus</i>	4 (3)	125
Uncharacterized protein (Ig-like domain-containing protein)	Nile tilapia		
A0A3B1JCF6_ASTMX	<i>Astyanax mexicanus</i>	6 (3)	123
IF rod domain-containing protein	Mexican tetra/blind cave fish		
*A0A3Q2YHX2_HIPCM	<i>Hippocampus comes</i>	3 (3)	122
Complement component 8 subunit beta	Tiger tail seahorse		
A0A3Q3IC70_MONAL	<i>Monopterus albus</i>	1 (1)	121
Ig-like domain-containing protein	Asian swamp eel		
*A0A0F8AI97_LARCR	<i>Larimichthys crocea</i>	2 (2)	121
Collagenase 3	Yellow croaker		
A0A6J2PEG5_COTGO	<i>Cottoperca gobio</i>	2 (2)	120
complement C5-like	Channel bull blenny		
A0A6A4TFM7_SCOMX	<i>Scophthalmus maximus</i>	3 (2)	119
Ig-like domain-containing protein	Turbot		
*A0A3Q0R4Z0_AMPPI	<i>Amphilophus citrinellus</i>	5 (3)	119
Complement component C9	Midas cichlid		
A0A6I9NNH1_9TELE	<i>Notothenia coriiceps</i>		
inter-alpha-trypsin inhibitor heavy chain H2	Black rockcod/Antarctic yellowbelly rockcod	3 (2)	118

A0A437D6V7_ORYJA	<i>Oryzias javanicus</i>	2 (2)	114
Chitinase	Javanese ricefish		
A0A3Q3EEY5_9LABR	<i>Labrus bergylta</i>	3 (3)	113
Fibrinogen C-terminal domain-containing protein	Ballan wrasse		
*A0A1S3SMN1_SALSA	<i>Salmo salar</i>	1 (1)	111
cathepsin L1-like	Atlantic salmon		
*A0A3Q3IZL2_MONAL	<i>Monopterus albus</i>	4 (3)	111
Uncharacterized protein	Asian swamp eel		
*A0A3P8YF02_ESOLU	<i>Esox Lucius</i>	6 (3)	109
Vitellogenin domain-containing protein	Northern pike		
*A0A3B3CJZ7_ORYME	<i>Oryzias melastigma</i>	3 (2)	109
Complement 4B (Chido blood group)	Marine medaka		
*A0A2U9CVZ8_SCOMX	<i>Scophthalmus maximus</i>	2 (2)	108
Putative complement component C8 gamma chain	Turbot		
A0A3Q3RJX0_9TELE	<i>Mastacembelus armatus</i>	2 (1)	108
Ig-like domain-containing protein	Zig-zag eel/Spiny eel		
*A0A3Q4HZS4_NEOBR	<i>Neolamprologus brichardi</i>	3 (3)	108
Uncharacterized protein (Ceruloplasmin)	Lyretail cichlid		
A0A6G1PYT4_9TELE	<i>Channa argus</i>		
Complement C5 C3 and PZP-like alpha-2-macroglobulin domain-containing protein 4	Northern snakehead	4 (3)	108
*A0A2U9AV20_SCOMX	<i>Scophthalmus maximus</i>	2 (2)	107
Prothrombin	Turbot		
A0A4W6EWH0_LATCA	<i>Lates calcarifer</i>	3 (3)	107
Peptidase S1 domain-containing protein	Barramundi/Asian sea bass		
*H3C6P0_TETNG	<i>Tetraodon nigroviridis</i>	2 (2)	106
Plasminogen	Green spotted puffer		
A0A3P8SDE5_AMPPE	<i>Amphiprion percula</i>	14 (2)	105
Serotransferrin	Orange clownfish		
A0A3B4BP10_PYGNA	<i>Pygocentrus nattereri</i>	10 (2)	105
Uncharacterized protein	Red-bellied piranha		
*A0A3B4TQB5_SERDU	<i>Seriola dumerili</i>	1 (1)	105
SERPIN domain-containing protein	Greater amberjack		
*D5A7I1_DICLA	<i>Dicentrarchus labrax</i>	4 (2)	104
Hemopexin	European bass		
*A0A2U9CU10_SCOMX	<i>Scophthalmus maximus</i>		
Putative insulin-like growth factor-binding protein complex acid labile subunit	Turbot	3 (3)	103
*A0A6J2PA80_COTGO	<i>Cottoperca gobio</i>	3 (3)	103
histone H2B 1/2-like	Channel bull blenny		
*A0A3P8SSL4_AMPPE	<i>Amphiprion percula</i>		
Uncharacterized protein (Ig-like domain-containing protein, Nattectin)	Orange clownfish	2 (2)	102
*G3NN36_GASAC	<i>Gasterosteus aculeatus</i>	4 (3)	99
Uncharacterized protein	Three-spined stickleback		
A0A4W4FLR8_ELEEL	<i>Electrophorus electricus</i>	3 (2)	99
Fibrinogen beta chain	Electric eel		
A0A671TDU8_SPAAU	<i>Sparus aurata</i>	2 (2)	97
Ig-like domain-containing protein	Gilt-head bream		
A0A6I9PPY0_9TELE	<i>Notothenia coriiceps</i>		
fibrinogen gamma chain	Black rockcod/Antarctic yellowbelly rockcod	2 (2)	96
A0A671TNW0_SPAAU	<i>Sparus aurata</i>	4 (3)	96
Histone H3	Gilt-head bream		
*A0A3B4XVK3_SERLL	<i>Seriola lalandi dorsalis</i>	2 (2)	96
Vitellogenin domain-containing protein	Yellowtail amberjack		
*A0A3Q3L1F9_9TELE	<i>Mastacembelus armatus</i>	1 (1)	95

Complement component 1, r subcomponent	Zig-zag eel/Spiny eel		
A0A1A8AN27_NOTFU	<i>Nothobranchius furzeri</i>	3 (3)	95
Fibrinogen, gamma polypeptide	turquoise killifish		
*A0A2D0QC28 ICTPU	<i>Ictalurus punctatus</i>	2 (2)	93
Ig heavy chain Mem5-like	Channel catfish		
A0A3P8R4C1_ASTCA	<i>Astatotilapia calliptera</i>		
Uncharacterized protein (Ig-like domain-containing protein)	Eastern happy/eastern river bream	6 (2)	96
A0A3B4H9E9_9CICH	<i>Pundamilia nyererei</i>	3 (2)	93
Ig-like domain-containing protein	Cichlid		
A0A3B4UNU3_SERDU	<i>Seriola dumerili</i>	4 (2)	93
Ig-like domain-containing protein	Greater amberjack		
*A0A060XWP2_ONCMY	<i>Oncorhynchus mykiss</i>		92
SERPIN domain-containing protein	Rainbow trout		
*A0A1A8CRV1_9TELE	<i>Nothobranchius kadleci</i>	8 (2)	91
Uncharacterized protein	Killifish		
*A0A2U9CFI3_SCOMX	<i>Scophthalmus maximus</i>	2 (2)	90
Putative sushi domain-containing protein 2 isoform 2	Turbot		
A0A5C6NS08_9TELE	<i>Takifugu flavidus</i>	5 (2)	90
Ig heavy chain V region VH558 A1/A4	Yellowbelly pufferfish		
*A0A4W4DXU4_ELEEL	<i>Electrophorus electricus</i>	3 (3)	89
14_3_3 domain-containing protein	Electric eel		
*A0A0F8B5M5_LARCR	<i>Larimichthys crocea</i>	1 (1)	88
Catechol O-methyltransferase domain-containing protein 1	Yellow croaker		
*A0A5N5KRW8_PANHP	<i>Pangasianodon hypophthalmus</i>	3 (3)	88
Uncharacterized protein (pleckstrin homology domain-containing family)	Iridescent shark		
*A0A5C6NRB2_9TELE	<i>Takifugu flavidus</i>	2 (2)	87
Apolipoprotein B-100	Yellowbelly pufferfish		
A0A2D0RGG9 ICTPU	<i>Ictalurus punctatus</i>	3 (2)	87
catenin beta-1 isoform X3	Channel catfish		
*A0A6I9P4Q9_9TELE	<i>Notothenia coriiceps</i>	1 (1)	86
apolipoprotein B-100-like	Black rockcod/Antarctic yellowbelly rockcod		
*A0A087XVJ8_POEFO	<i>Poecilia formosa</i>	2 (1)	86
Uncharacterized protein (IGv domain-containing protein)	Amazon molly		
*H1AB41_PLASA	<i>Platichthys stellatus</i>	4 (2)	85
Lysozyme	Starry flounder		
*A0A4P8JEC9_9TELE	<i>Lateolabrax maculatus</i>	2 (2)	84
Apolipoprotein Ba	Spotted sea bass		
A0A3Q4ACH4_MOLML	<i>Mola mola</i>	2 (2)	84
Inter-alpha-trypsin inhibitor heavy chain 3	Ocean sunfish		
*A0A484CC61_PERFV	<i>Perca flavescens</i>	3 (1)	84
Uncharacterized protein (Hyaluronan-binding protein 2)	Yellow perch		
A0A060Z3N3_ONCMY	<i>Oncorhynchus mykiss</i>	3 (2)	86
Ig-like domain-containing protein	Rainbow trout		
*A0A3B4ZU87_9TELE	<i>Stegastes partitus</i>	3 (2)	83
Uncharacterized protein (complement factor H-like)	Bicolour damselfish		
A0A3B3QDE5_9TELE	<i>Paramormyrops kingsleyae</i>	2 (1)	83
Ig-like domain-containing protein	Elephantfish		
A0A3B3CFL8_ORYME	<i>Oryzias melastigma</i>	5 (2)	83
Ig-like domain-containing protein	Marine medaka		
*A0A3Q3W6Q7_MOLML	<i>Mola mola</i>	2 (2)	82
Sushi domain containing 2	Ocean sunfish		
A0A4W5L5T6_9TELE	<i>Hucho hucho</i>	3 (2)	82
Thioredoxin	Danube salmon		
*G1DHP8_GOBRA	<i>Gobiocypris rarus</i>	2 (2)	81

Vitellogenin (Fragment)	Rare gudgeon/rare minnow		
*A0A3B3QP35_9TELE	<i>Paramormyrops kingsleyae</i>	3 (2)	80
Uncharacterized protein	Elephantfish		
*A0A3B4Z082_9TELE	<i>Stegastes partitus</i>	2 (2)	80
Uncharacterized protein (complement C6)	Bicolour damselfish		
A0A669CCK4_ORENI	<i>Oreochromis niloticus</i>	6 (2)	80
Uncharacterized protein (Ig-like domain-containing protein)	Nile tilapia		
*A0A484C6M0_PERFV	<i>Perca flavescens</i>	1 (1)	80
Uncharacterized protein	Yellow perch		
*A0A3P8U2B4_AMPPE	<i>Amphiprion percula</i>	4 (2)	80
Keratin 98	Orange clownfish		
*A0A060WHH8_ONCMY	<i>Oncorhynchus mykiss</i>	2 (2)	79
Junction plakoglobin	Rainbow trout		
A0A3B4ULY5_SERDU	<i>Seriola dumerili</i>	4 (2)	78
Ig-like domain-containing protein	Greater amberjack		
H3C0U1_TETNG	<i>Tetraodon nigroviridis</i>	3 (2)	77
Ig-like domain-containing protein	Green spotted puffer		
A0A087X4F8_POEFO	<i>Poecilia formosa</i>	1 (1)	77
Uncharacterized protein (Ig-like domain-containing protein)	Amazon molly		
A0A3P9IRN4_ORYLA	<i>Oryzias latipes</i>	2 (2)	77
Ig-like domain-containing protein	Medaka/Japanese rice fish		
A0A060W543_ONCMY	<i>Oncorhynchus mykiss</i>	2 (2)	77
Histone H2A	Rainbow trout		
A0A3B4UFJ1_SERDU	<i>Seriola dumerili</i>	2 (1)	75
Ig-like domain-containing protein	Greater amberjack		
A0A0F8ABH4_LARCR	<i>Larimichthys crocea</i>	5 (1)	75
Granzyme B(G,H)	Yellow croaker		
*A0A3B4UPX8_SERDU	<i>Seriola dumerili</i>	1 (1)	74
Zona pellucida sperm-binding protein 3	Greater amberjack		
*A0A3P8U813_AMPPE	<i>Amphiprion percula</i>	2 (2)	73
Si:ch1073-416d2.3	Orange clownfish		
*A0A3Q1KAD2_ANATE	<i>Anabas testudineus</i>	2 (2)	72
SERPIN domain-containing protein	Climbing perch		
*A0A4W5RID4_9TELE	<i>Hucho hucho</i>	2 (1)	71
RRM domain-containing protein	Danube salmon		
*A0A3Q2QNZ9_FUNHE	<i>Fundulus heteroclitus</i>	2 (2)	71
Uncharacterized protein (Sushi domain containing 2)	Atlantic killifish, mud minnow		
*A0A4W5LQ29_9TELE	<i>Hucho hucho</i>	8 (2)	70
ATP-synt ab_N domain-containing protein	Danube salmon		
A0A3Q2PS35_FUNHE	<i>Fundulus heteroclitus</i>	5 (2)	70
Ig-like domain-containing protein	Atlantic killifish, mud minnow		
*A0A6G1QID3_9TELE	<i>Channa argus</i>	2 (2)	70
Complement component C6	Northern snakehead		
*A0A3B3X986_9TELE	<i>Poecilia Mexicana</i>	1 (1)	70
Uncharacterized protein (F-BAR domain-containing protein)	Atlantic (shortfin) molly		
*A0A498LNY2_LABRO	<i>Labeo rohita</i>	6 (2)	70
Retrotransposon-derived PEG10	Rohu		
*A0A6G1PD67_9TELE	<i>Channa argus</i>	2 (2)	70
Apoptosis-stimulating of p53 protein 2 Bcl2-binding protein	Northern snakehead		
*A0A1S3L2W1_SALSA	<i>Salmo salar</i>	5 (2)	70
FH2 domain-containing protein 1-like	Atlantic salmon		
A0A3B3HM39_ORYLA	<i>Oryzias latipes</i>	1 (1)	69
Ig-like domain-containing protein	Medaka/Japanese rice fish		
*A0A3Q1HK94_ANATE	<i>Anabas testudineus</i>	6 (2)	69
Protein-tyrosine-phosphatase	Climbing perch		
A0A3Q3JUN7_MONAL	<i>Monopterus albus</i>	2 (2)	68
IF rod domain-containing protein	Asian swamp eel		

*A0A671X983_SPAAU	<i>Sparus aurata</i>		
Uncharacterized protein (Early endosome antigen 1, FYVE-type domain-containing protein)	Gilt-head bream	3 (2)	68
*A0A3B3DTR8_ORYME	<i>Oryzias melastigma</i>		
Uncharacterized protein	Marine medaka	3 (2)	68
*A0A3Q3XI23_MOLML	<i>Mola mola</i>		
Zgc:112265	Ocean sunfish	3 (2)	67
A0A671YT10_SPAAU	<i>Sparus aurata</i>		
Uncharacterized protein (Immunoglobulin like and fibronectin type III domain containing 1, tandem duplicate 2)	Gilt-head bream	2 (2)	67
*A0A3B5ACM2_9TELE	<i>Stegastes partitus</i>		
Uncharacterized protein	Bicolour damselfish	6 (2)	66
A0A3P9H0Y9_ORYLA	<i>Oryzias latipes</i>		
Ig-like domain-containing protein	Medaka/Japanese rice fish	2 (2)	65
*A0A5C6N3H2_9TELE	<i>Takifugu flavidus</i>		
Keratin, type I cytoskeletal 18	Yellowbelly pufferfish	4 (2)	65
*A0A3B5L5A5_9TELE	<i>Xiphophorus couchianus</i>		
Thyroid hormone receptor interactor 11	Monterrey platyfish	3 (2)	65
*Q2PZ29_SOLSE	<i>Solea senegalensis</i>		
Lysozyme	Senegalese sole	2 (1)	65
A0A667YBU1_9TELE	<i>Myripristis murdjan</i>		
Ig-like domain-containing protein	Blacktipped soldierfish	5 (2)	65
*A0A672GWK0_SALFA	<i>Salarias fasciatus</i>		
Uncharacterized protein (Complement factor B-like)	Lawnmower blenny	2 (2)	64
*A0A3B4CEW8_PYGNA	<i>Pygocentrus nattereri</i>		
Uncharacterized protein (Roundabout-like axon guidance receptor protein 2)	Red-bellied piranha	2 (2)	64
*A0A3B4EX20_9CICH	<i>Pundamilia nyererei</i>		
Uncharacterized protein (Apolipoprotein M)	Cichlid	6 (2)	64
*A0A2I4BMF1_9TELE	<i>Austrofundulus limnaeus</i>		
protein Z-dependent protease inhibitor-like	Killifish	1 (1)	63
*A0A3Q3EPX4_9LABR	<i>Labrus bergylta</i>		
Vitellogenin domain-containing protein	Ballan wrasse	2 (2)	62
*A0A3B4T5U4_SERDU	<i>Seriola dumerili</i>		
Uncharacterized protein (Myosin phosphatase Rho interacting protein)	Greater amberjack	3 (2)	62
A0A3B3T2D8_9TELE	<i>Paramormyrops kingsleyae</i>		
Ig-like domain-containing protein	Elephantfish	1 (1)	62
*A0A3Q1FWV1_9TELE	<i>Acanthochromis polyacanthus</i>		
Multidrug and toxin extrusion protein	Spiny chromis damselfish	2 (2)	62
A0A3B4YHZ5_SERLL	<i>Seriola lalandi dorsalis</i>		
IGv domain-containing protein	Yellowtail amberjack	1 (1)	61
*R4I5B0_EPICO	<i>Epinephelus coioides</i>		
Immunoglobulin light chain	Orange-spotted grouper	3 (2)	61
*A0A3Q0R568_AMPCI	<i>Amphilophus citrinellus</i>		
FH2 domain containing 4	Midas cichlid	3 (2)	61
A0A3B4WXW5_SERLL	<i>Seriola lalandi dorsalis</i>		
Ig-like domain-containing protein	Yellowtail amberjack	2 (2)	60
G3PK20_GASAC	<i>Gasterosteus aculeatus</i>		
Serotransferrin	Three-spined stickleback	3 (2)	60
*A0A484DB45_PERFV	<i>Perca flavescens</i>		
Uncharacterized protein (Pentaxin)	Yellow perch	1 (1)	60
*A0A671SV95_9TELE	<i>Sinocyclocheilus anshuiensis</i>		
FERM domain-containing protein	Sinocyclocheilus cavefish (Cyprinoid)	2 (2)	60
A0A023REA6_9TELE	<i>Menidia estor</i>		
Elongation factor 1- α	Pike silverside	1 (1)	60

*A0A6J2PC09_COTGO <i>nesprin-2</i>	<i>Cottoperca gobio</i>	2 (2)	60
*A0A0S7MGP3_9TELE ZN287 (Fragment)	Channel bull blenny <i>Poeciliopsis prolifica</i>	3 (2)	59
*A0A3Q3VSX4_MOLML Uncharacterized protein	Blackstripe livebearer <i>Mola mola</i>	1 (1)	59
*A0A553Q8B1_9TELE Uncharacterized protein	Ocean sunfish <i>Danionella translucida</i>	3 (2)	58
*A0A0P7TM62_SCLFO Keratin, type I cytoskeletal 18-like	Micro glassfish (Cyprinid) <i>Scleropages formosus</i>	1 (1)	58
*A0A060XKV1_ONCMY [Histone H3]-trimethyl-L-lysine(9) demethylase	Asian arowana <i>Oncorhynchus mykiss</i>	3 (2)	58
*E7F6Y7_DANRE DNA polymerase kappa	Rainbow trout <i>Danio rerio</i>	4 (2)	58
*F8W5U5_DANRE Centrosomal protein of 290 kDa	Zebrafish <i>Danio rerio</i>	2 (2)	58
*A0A2U9CTT6_SCOMX Putative utrophin	Zebrafish <i>Scophthalmus maximus</i>	7 (2)	57
*A0A3B3BVC4_ORYME Uncharacterized protein	Turbot <i>Oryzias melastigma</i>	2 (2)	57
*A0A3B4UZF1_SERDU [Histone H3]-lysine(4) N-trimethyltransferase	Marine medaka <i>Seriola dumerili</i>	1 (1)	57
A0A060VW86_ONCMY Uncharacterized protein (Tubulin alpha, tubulin domain containing)	Greater amberjack <i>Oncorhynchus mykiss</i>	1 (1)	56
*A0A671TLU7_SPAAU Reverse transcriptase	Rainbow trout <i>Sparus aurata</i>	3 (2)	56
A0A3Q4H8B0_NEOBR Ig-like domain-containing protein	Gilt-head bream <i>Neolamprologus brichardi</i>	1 (1)	56
*A0A0U2ERZ3_CORCL Glyceraldehyde 3-phosphate dehydrogenase	Lyretail cichlid <i>Coregonus clupeaformis</i>	6 (1)	56
*A0A0R4IVM1_DANRE LSM14A mRNA-processing body assembly factor b	Lake whitefish <i>Danio rerio</i>	11 (2)	55
*A0A3P8VC95_CYNSE Uncharacterized protein	Zebrafish <i>Cynoglossus semilaevis</i>	1 (1)	54
*Q9DFN6_GILMI Glyceraldehyde-3-phosphate dehydrogenase	Tongue sole <i>Gillichthys mirabilis</i>	1 (1)	54
*A0A3B3BWJ2_ORYME Uncharacterized protein	<i>Oryzias melastigma</i>	2 (2)	54
*A0A6A4SGZ4_SCOMX C1q domain-containing protein	Marine medaka <i>Scophthalmus maximus</i>	1 (1)	54
*A0A3B4EJ56_PYGNA von Willebrand factor	Turbot <i>Pygocentrus nattereri</i>	2 (2)	54
*A0A1S3RE28_SALSA uncharacterized protein LOC106602330 isoform X1	Red-bellied piranha <i>Salmo salar</i>	1 (1)	53
*A0A2I4CMN8_9TELE titin-like	Atlantic salmon <i>Austrofundulus limnaeus</i>	2 (2)	53
	Killifish		

[†] Ions score is $-10 \cdot \log(P)$, where P is the probability that the observed match is a random event. Individual ions scores > 53 indicate identity or extensive homology ($p < 0.05$). Protein scores are derived from ions scores as a non-probabilistic basis for ranking protein hits.




Figure 2. The proteome and citrullinome of halibut serum-EVs. Silver-stained gels for: (A) total protein cargo in EVs and (B) F95 enriched (deiminated/citrullinated) proteins from EVs. The protein standard (std) is indicated in kilodaltons (kDa). (C) Venn diagram shows the number of candidate protein hits identified as cargo in total serum EVs (“The serum EV proteome”) as well as deiminated protein hits in EV cargo (the serum “EV citrullinome”).

2.3. Complement Component C3, C4 and Pentraxin-Like Protein verified in Halibut EVs and F95 Enriched EV Protein Cargo Fractions using Western Blotting

Three candidate proteins which were identified as part of EV total protein cargo by LC-MS/MS, namely complement component C3, C4 and pentraxin-like protein, were further assessed by western blotting in halibut serum-EVs (Figure 3A–C). Both total EV protein cargo as well as the F95 enriched protein cargo were assessed, using halibut-specific C3, C4 and pentraxin-like protein antibodies, respectively, which had previously been generated and validated in our laboratories [13,42]. Here, complement component C3 was verified to be present in total EV protein cargo, where it was strongly detected by western blotting, as well as at lower levels in the deiminated (F95-enriched) protein cargo (Figure 3A). This confirmed the hits identified by the LC-MS/MS analysis, showing that C3 is exported in EVs both in normal and deiminated form (Table 1; Table 2). Complement component C4 was also confirmed to be exported in total EV cargo by western blotting, albeit at lower levels than C3, in accordance with the LC-MS/MS findings which identified C4 as a hit in total EV cargo. C4 was seen only at very low levels in deiminated form in the F95-enriched EV fraction by Western blotting (Figure 3B), and was not identified as part of the F95-enriched cargo by LC-MS/MS. Pentraxin-like protein was strongly detected in total EV protein cargo by western blotting, but not in the F95-enriched EV protein fractions (Figure 3C), in accordance with the results from the LC-MS/MS analysis, which only detected pentraxin in total EV cargo (Table 2).






Figure 3. Complement component C3, C4 and pentraxin-like protein in halibut EVs and F95 enriched EV fractions. Western blotting showing (A) complement component C3 detection in total protein cargo of halibut serum-EVs (“EVs”) and in F95-enriched protein fractions from serum-EVs (“EVs_F95”), C3 α - and β -chains, as well as α -fragment (α -f) are indicated; (B) complement component C4 detection in total protein cargo of serum-EVs (“EVs”) and lower detection observed in F95-enriched EV protein fractions (“EVs_F95”), C4 α -, β - and γ -chains are indicated; (C) pentraxin-like protein detection in total EV protein cargo (“EVs”), which was not detected in the F95-enriched EV protein fractions (“EVs_F95”).

2.4. Protein–Protein Interaction Network Analysis for Halibut Serum-EV Protein Cargo: Deiminated and Total Protein Cargo

2.4.1. Protein Interaction Networks Enriched for Halibut Serum-EV Deiminated/Citrullinated Protein Cargo

For the generation of protein–protein interaction networks to further understanding of putative protein pathways regulated by deimination, deiminated (F95-enriched) protein hits from halibut EVs were assessed by STRING analysis. The protein hits were assessed using the general teleost STRING database, selecting the zebrafish (*Danio rerio*) database as a model database, as no specific database for halibut is available in STRING and zebrafish showed the highest identity with the teleost protein hits identified as deiminated in halibut serum-EVs. The protein–protein interaction networks showed a PPI enrichment p -value of 5.15×10^{-5} , indicating significantly more interactions than expected from a random set of proteins (Figure 4).






Figure 4. (A) Protein interaction networks for deiminated proteins in halibut EVs. Local network clusters and UniProt keywords are indicated by the colour coded nodes. See colour key for nodes and interaction networks in the figure. (B) Reactome protein interaction networks for deiminated proteins in halibut EVs. Reactome pathways are indicated by the coloured nodes, as shown in the figure. (C,D) PFAM and SMART protein interaction networks for deiminated proteins in halibut EVs. PFAM and SMART protein domains are indicated by the coloured nodes, see colour code in the figure. (E) InterPro protein interaction networks for deiminated proteins in halibut EVs. InterPro protein domains and features are indicated by the coloured nodes; see colour code in the figure.

Local network clusters enriched in deiminated proteins in EVs included: Histone H3/CENP-A, core histone H2A/H2B/H3/H4 network, post-translational protein phosphorylation and the regulation of IGF transport (Figure 4A).

UniProt keywords for deiminated proteins identified in serum-EVs included methylation, cytoskeleton, disulphide bond, cytoplasm and signalling (Figure 4A).

Reactome pathways enriched in deiminated proteins in the serum EVs included GRB2:SOS linkage to MAPK signalling for integrins, p130Cas linkage to MAPK signalling for integrins, MAP2K and MAPK activation, integrin signalling, the initial triggering of complement, L1CAM interactions, post-translational protein phosphorylation, the regulation of actin dynamics for phagocytic cup formation, the regulation of IGF transport, platelet degranulation, integrin cell surface interactions, cell junction organisation, clathrin-mediated endocytosis, VEGFA-VEGFR2 pathway, extracellular matrix organisation, developmental biology, innate immune system and neutrophil degranulation (Figure 4B).


PFAM protein domains for deiminated proteins identified in the serum EVs included alpha-macroglobulin thiolester bond-forming region, alpha-2-macroglobulin family N-terminal region, MG2 domain, alpha-2 macroglobulin family, a-macroglobulin complement component, a-macroglobulin receptor, UNC-6/NTR/C345C module, core histone H2A/H2B/H3/H4 and trypsin (Figure 4C).


SMART protein domains for deiminated EV proteins included alpha-macroglobulin family, alpha-2-macroglobulin, a-macroglobulin receptor, kazal type serine protease inhibitors, domains found in plexins, semaphorins and integrins and trypsin-like serine protease (Figure 4D).

Protein domains and features (InterPro) for deiminated proteins in serum-EVs included macroglobulin domain MG4 and MG3, tissue inhibitor of metalloproteinases-like, OB-fold, netrin module, non-TIMP type, netrin domain, PSI domain, peptidase S1A, chymotrypsin family, peptidase S1 PA clan, serine proteases trypsin family, histidine active site and serine active site (Figure 4E).

2.4.2. Protein Interaction Networks enriched for Halibut Serum-EV Total Protein Cargo

The same approach for the generation of protein–protein interaction networks, selecting the zebrafish (*D. rerio*) STRING database as a representative database for teleost fish, was also applied for total protein EV cargo identified in halibut, showing a PPI enrichment p -value: $<1.0 \times 10^{-16}$ for the protein networks generated, indicating significantly more interactions than expected from a random set of proteins (Figure 5).



Reactome-3**Reactome-3:**

- Binding and uptake of ligands by scavenger receptors
- Platelet degranulation
- Collagen degradation
- Integrin cell surface interactions
- Metabolism of vitamins and cofactors
- Toll-like receptor cascades
- Platelet activation, signalling and aggregation
- Clathrin-mediated endocytosis
- Peptide ligand-binding receptors
- Extracellular matrix organisation


Reactome-4**Reactome-4:**

- Innate immune system
- G alpha signalling events
- Hemostasis
- Developmental biology
- Neutrophil degranulation
- GPCR downstream signalling
- Post-translational protein modification
- Signal transduction
- Metabolism of proteins

C.**UniProt****UniProt:**

- Kringle
- Methylation
- Nucleosome core
- Serine protease
- Secreted
- Chromosome
- Protease
- Disulphide bond
- Signal

D.


**PFAM-1:**

- Fibrinogen alpha/beta chain family
- Anaphylatoxin-like domain
- Vault protein inter-alpha-trypsin domain
- MG2 domain
- Alpha-2-macroglobulin family
- Alpha-2-macroglobulin complement component
- Lipoprotein amino terminal region
- UNC-6/NTR/C345C module
- Kringle domain
- Domain of unknown function (DUF1943)

**PFAM-2:**

- vWf type D domain
- Hemopexin
- vWf type A domain
- FYVE zinc finger
- Fibrinogen beta, gamma chains, C-terminal globular domain
- Trypsin-like peptidase domain
- CUB domain
- Core histone H2A/H2B/H3/H4
- Trypsin
- vWf type A domain

E.

**SMART-1:**

- Fibrinogen alpha/beta chain family
- Anaphylatoxin homologous domain
- Vault protein inter-alpha-trypsin domain
- Kringle domain
- Lipoprotein N-terminal domain
- Netrin C-terminal domain
- Large open beta-sheet protein family
- Alpha-2-macroglobulin family
- Alpha-2-macroglobulin
- Alpha-2-macroglobulin receptor

**SMART-2:**

- vWf type D domain
- Hemopexin-like repeats
- Protein present in Fab1, YOTB, Vac1 and EEA1
- Fibrinogen related domains (FReDs)
- Kazal type serine protease inhibitors
- Domain first found in C1r, C1s, uEGF and bone morphogenesis
- vWf type A domain
- Trypsin-like serine protease




Figure 5. (A) Protein interaction networks for total protein cargo in halibut EVs, showing local network clusters. The coloured nodes indicate the different networks, respectively. (B) Reactome protein interaction networks for total proteins in halibut EV cargo, showing reactome pathways. Specific reactome pathways are indicated by the coloured nodes, respectively. (C) UniProt protein interaction networks for total proteins in halibut EV cargo, showing UniProt keywords. UniProt keywords are indicated by the coloured nodes, respectively. (D) PFAM protein interaction networks for total proteins in halibut EV cargo. The specific PFAM protein domains are indicated by

the coloured nodes, respectively. (E) Protein interaction networks for total proteins in halibut EVs, showing SMART protein domains. The specific SMART protein domains are indicated by the coloured nodes, respectively. (F) InterPro protein interaction networks for total proteins in halibut EVs. The specific protein domains and features (InterPro) are indicated by the coloured nodes, respectively.

Local network clusters for total EV protein content included fibrinogen family, fibrinolysis, common pathway of fibrin clot formation, clotting cascade, ApoM domain, selenoprotein P, histone H3/CENP-A, histone H4, terminal pathway of complement, alternative complement activation, MG2 domain, terminal complement pathway, lectin pathway, plakophilin/delta catenin desmosomal, regulation of IGF transport, adherens junctions interactions, MHC class II antigen, MHC class II antigen presentation, post-translational protein phosphorylation, Histone H4, and core histone H2A/H2B/H3/H4 (Figure 5A).

Reactome pathways for total EV protein cargo included LDL remodelling, plasma lipoprotein assembly, remodelling and clearance, innate immune system, Toll-like receptor cascades, neutrophil degranulation, the regulation of complement cascade, terminal complement pathway, the activation of C3 and C5, the initial triggering of complement, platelet degranulation, platelet activation, GRB2:SOS linkage to MAPK, integrin signalling, integrin cell surface interactions, p130Cas linkage to MAPK signalling for integrins, MAP2K and MAPK activation, the activation of matrix metalloproteinases, chylomicron assembly, the common pathway of fibrin clot formation, the intrinsic pathway of fibrin clot formation, the formation of fibrin clot, clotting cascade, platelet aggregation (plug formation), the formation of the cornified envelope, the regulation of IGF transport, the binding and uptake of ligands by scavenger receptors, collagen degradation, the metabolism of vitamins and cofactors, retinoid metabolism and transport, clathrin-mediated endocytosis, peptide ligand-binding receptors, extracellular matrix organisation, G alpha signalling events, hemostasis, signalling and aggregation, developmental biology, GPCR downstream signalling, post-translational protein modification, signal transduction, metabolism of proteins (Figure 5B).

UniProt keywords for total EV protein content included methylation, kringle, nucleosome core, serine protease, secreted, chromosome, protease, disulphide bond, signalling (Figure 5C).

PFAM protein domains for total EV protein cargo included fibrinogen alpha/beta chain family, anaphylatoxin-like domain, vault protein inter-alpha-trypsin domain, MG2 domain, alpha-2-macroglobulin family, alpha-2-macroglobulin complement component, lipoprotein amino terminal region, UNC-6/NTR/C345C module, kringle domain, domain of unknown function (DUF1943), vWf type A domain, vWf type D domain, hemopexin, FYVE zinc finger, fibrinogen beta, gamma chains, C-terminal globular domain, trypsin-like peptidase domain, CUB domain, core histone H2A/H2B/H3/H4, and trypsin (Figure 5D).

SMART protein domains for total EV protein cargo included fibrinogen alpha/beta chain family, anaphylatoxin homologous domain, vault protein inter-alpha-trypsin domain, kringle domain, lipoprotein N-terminal domain, netrin C-terminal domain, large open beta-sheet protein family, alpha-2-macroglobulin family, alpha-2-macroglobulin receptor, vWf type A domain, vWf type D domain, hemopexin-like repeats, protein present in Fab1, YOTB, Vac1 and EEA1, fibrinogen related domains (FReDs), kazal type serine protease inhibitors, domain first found in C1r, C1s, uEGF and bone morphogenesis, and trypsin-like serine protease (Figure 5E).

Protein domains and features (InterPro) identified for total EV cargo included fibrinogen alpha/beta/gamma chain coiled-coil domain, complement C3/4/5, MG1 domain, anaphylatoxin, anaphylatoxin/fibulin, complement system, macroglobulin domain, alpha-macroglobulin TED domain, alpha-2-macroglobulin, VIT domain, kringle superfamily, terpenoid cyclases/protein prenyltransferase alpha-alpha toroid, zinc finger,

FYVE related, FYVE zinc finger, immunoglobulin-like fold, lipid transport protein, beta-sheet shell, lipovitellin-phosvitin complex, superhelical domain, lipid transport protein, netrin domain, netrin module non-TIMP type, vWf type A and vWf type D domain, tissue inhibitor of metalloproteinases-like OB-fold, histone H2A/H2B/H3, histone-fold, peptidase S1, PA clam, fibrinogen alpha/beta/gamma chain C-terminal globular domain, fibrinogen-like C-terminal, serine proteases, trypsin family serine active site and histidine active site, sushi/SCR/CCP superfamily, and peptidase S1A chymotrypsin family (Figure 5F).

3. Discussion

This is the first study to assess EV profile signatures in halibut biofluids, identifying both total serum-EV protein cargo as well as deiminated protein cargo in serum-EVs. The size profiling of halibut serum-EVs by NTA showed vesicles in the range of 50–600 nm, which indicates a higher amount of larger EVs compared with human EVs, which typically fall in the size range of 30–300 nm. In comparison, while few teleost fish have been profiled for EVs, cod (*Gadus morhua*), serum-EVs were found to be in the size range of mainly 50–300 nm [33,38], while cod mucus-EVs are in the size range of 50–500 nm [32]. In other taxa across the phylogeny tree, differences in plasma or serum EV size profiles have indeed been reported. In elasmobranchs (nurse shark *Ginglymostoma cirratum*) a higher abundance of small EVs in the 10–200 nm size range was observed [14]; in a group of eight pelagic seabird species, some species-specific differences were reported showing plasma-EVs at 50–200 nm size range for some birds and others showing larger EVs at 250–500 nm [37], while in reptile (alligator—*Alligator mississippiensis*), plasma EVs were in the size range of 50–400 nm [16]. In llama (*Lama glama*), plasma-EVs were reported at 40–400 nm [34], while *Bos taurus* plasma-EV showed size profiles of 70–500 nm [35]. Naked mole-rat (*Heterocephalus glaber*) plasma shows similar EV size profiles as human plasma at 50–300 nm [38], as does rat (*Rattus norvegicus*) plasma at 50–250 nm [43]. In sea mammals, such as pinnipeds and cetaceans, serum-EVs were observed at 50–600 nm in seals [40], similar to as observed in halibut in the current study. In four species of whale, EV profiles were seen in the ranges of 50–500 (minke whale *Balaenoptera acutorostrata*), 50–400 (fin whale *Balaenoptera physalus*), 80–300 (humpback whale *Megaptera novaeangliae*) and 90–300 nm (Cuvier's beaked whale *Ziphius cavirostris*), respectively, while orca serum-EVs (*Orcinus orca*; dolphin family) were reported at 30–500 nm [39]. Reports of EV profiling of haemolymph from species lower in the phylogeny tree include Crustacea (lobster *Homarus americanus*) with EVs in the 10–500 nm size range (with the majority of EVs being small in the 22–115 nm size range) [22]; Mollusca haemolymph EVs at 50–300 nm (blue mussel, *Mytilus edulis*), 30–300 nm (soft shell clam *Mya arenaria*), 90–500 nm (Eastern oyster *Crassostrea virginica*) and 20–300 nm (Atlantic jackknife clam *Ensis leei*), respectively [24]; Arthropoda (horseshoe crab *Limulus polyphemus*) EVs at 20–400 nm (with the majority of EVs falling within 40–123 nm) [23]. In the protozoa *Giardia intestinalis*, two distinct size populations of EVs have been described (20–80 nm and 100–400 nm, respectively), which display different functions in host–pathogen interactions [21]. In Gram-negative and Gram-positive bacteria, with EV profiles described at 10–600 nm and 60–400 nm, respectively, EV profiles were shown to change in response to drug-treatment both with respect to size profile and EV cargo content [19,44]. This does indicate that EV size profiles differ between taxa and this may, amongst others, also have effects on EV cargo content, including proteomic, post-translationally modified proteomic cargo, as well as other genomic and non-coding RNA and mitochondrial-derived cargo [45]. Indeed, in teleost, it has been reported that changes in EV numbers and EV deimination protein and microRNA cargo can be a biomarker for environmental temperature factors [33] and, in response to other stressors, teleost plasma EVs have been found enriched with Hsp70 [46] and selected micro-RNAs [47]. In human parasitic disease, EV profiles can also be indicative of infection status [48]. Therefore, the characterisation of EVs across a wide range of

taxa further highlights their potential for biomarker application or “EV-fingerprinting” for the assessment of animal health.

Analysing both whole proteomic and the deiminated protein content of halibut serum-EVs in the current study, some differences were found in protein-interaction pathways, while overall both the whole proteome and the EV-citrullinome involved a number of immune, metabolic and gene regulatory pathways.

When assessing protein-protein interaction networks for EVs enriched in deiminated proteins, these related to local network clusters for deiminated proteins in serum-EVs included histone H3/CENP-A, core histone H2A/H2B/H3/H4 network, post-translational protein phosphorylation and the regulation of IGF transport. In relation to such networks, UniProt keywords for deiminated proteins identified in serum-EVs included methylation, cytoskeleton, disulphide bond, cytoplasm and signalling. Reactome pathways enriched in deiminated proteins in the serum EVs included GRB2:SOS linkage to MAPK signalling for integrins, p130Cas linkage to MAPK signalling for integrins, MAP2K and MAPK activation, integrin signalling, initial triggering of complement, L1CAM interactions, post-translational protein phosphorylation, regulation of actin dynamics for phagocytic cup formation, regulation of IGF transport, platelet degranulation, integrin cell surface interactions, cell junction organisation, clathrin-mediated endocytosis, VEGFA-VEGFR2 pathway, extracellular matrix organisation, developmental biology, innate immune system and neutrophil degranulation. Correspondingly, PFAM protein domains for deiminated proteins identified in the serum EVs included alpha-macro-globulin tie-ester bond-forming region, alpha-2-macroglobulin family N-terminal region, MG2 domain, alpha-2 macroglobulin family, alpha-macroglobulin complement component, alpha-macroglobulin receptor, UNC-6/NTR/C345C module, core histone H2A/H2B/H3/H4 and trypsin. SMART protein domains for deiminated EV proteins included alpha-macroglobulin family, alpha-2-macroglobulin, alpha-macroglobulin receptor, kazal-type serine protease inhibitors, domains found in plexins, semaphorins and integrins and trypsin-like serine. Protein domains and features (InterPro) for deiminated proteins in serum-EVs included macroglobulin domain MG4 and MG3, the tissue inhibitor of metalloproteinases-like OB-fold, netrin module, non-TIMP type, netrin domain, PSI domain, peptidase S1A, chymotrypsin family, peptidase S1 PA clan, serine proteases trypsin family, histidine active site and serine active site.

In comparison with deiminated EV protein content, more pathways were revealed for serum-EV total protein content, as would be expected due to only some of the proteins in the EV cargo being candidates for post-translational deimination and exported in EVs in deiminated form. Assessing protein interaction networks for total protein EV content showed local network clusters for fibrinogen family, fibrinolysis, the common pathway of fibrin clot formation, clotting cascade, ApoM domain, selenoprotein P, histone H3/CENP-A, histone H4, the terminal pathway of complement, alternative complement activation, MG2 domain, terminal complement pathway, lectin pathway, plakophilin/delta catenin desmosomal, the regulation of IGF transport, adherens junctions interactions, MHC class II antigen, MHC class II antigen presentation, post-translational protein phosphorylation, Histone H4, and core histone H2A/H2B/H3/H4.

The reactome pathways for total EV protein cargo included LDL remodelling, plasma lipoprotein assembly, remodelling and clearance, innate immune system, Toll-like receptor cascades, neutrophil degranulation, the regulation of the complement cascade, the terminal complement pathway, the activation of C3 and C5, the initial triggering of complement, platelet degranulation, platelet activation, GRB2:SOS linkage to MAPK, integrin signalling, integrin cell surface interactions, p130Cas linkage to MAPK signalling for integrins, MAP2K and MAPK activation, the activation of matrix metalloproteinases, chylomicron assembly, the common pathway of fibrin clot formation, the intrinsic pathway of fibrin clot formation, the formation of fibrin clot, clotting cascade,

platelet aggregation (plug formation), the formation of the cornified envelope, the regulation of IGF transport, the binding and uptake of ligands by scavenger receptors, collagen degradation, the metabolism of vitamins and cofactors, retinoid metabolism and transport, clathrin-mediated endocytosis, peptide ligand-binding receptors, extracellular matrix organisation, G alpha signalling events, hemostasis, signalling and aggregation, developmental biology, GPCR downstream signalling, post-translational protein modification, signal transduction, and the metabolism of proteins.

UniProt keywords for total EV protein content included methylation, kringle, nucleosome core, serine protease, secreted, chromosome, protease, disulphide bond, signalling.

PFAM protein domains for total EV protein cargo included fibrinogen alpha/beta chain family, anaphylatoxin-like domain, vault protein inter-alpha-trypsin domain, MG2 domain, alpha-2-macroglobulin family, alpha-2-macroglobulin complement component, lipoprotein amino terminal region, UNC-6/NTR/C345C module, kringle domain, the domain of unknown function (DUF1943), vWf type A domain, vWf type D domain, hemopexin, FYVE zinc finger, fibrinogen beta, gamma chains, C-terminal globular domain, trypsin-like peptidase domain, CUB domain, core histone H2A/H2B/H3/H4, and trypsin.

SMART protein domains for total EV protein cargo included fibrinogen alpha/beta chain family, anaphylatoxin homologous domain, vault protein inter-alpha-trypsin domain, kringle domain, lipoprotein N-terminal domain, netrin C-terminal domain, large open beta-sheet protein family, alpha-2-macroglobulin family, alpha-2-macroglobulin receptor, vWf type A domain, vWf type D domain, hemopexin-like repeats, protein present in Fab1, YOTB, Vac1 and EEA1, fibrinogen related domains (FReDs), kazal type serine protease inhibitors, domain first found in C1r, C1s, uEGF and bone morphogenesis, and trypsin-like serine protease.

Protein domains and features (InterPro) identified for total EV protein cargo included fibrinogen alpha/beta/gamma chain coiled-coil domain, complement C3/4/5, MG1 domain, anaphylatoxin, anaphylatoxin/fibulin, complement system, macroglobulin domain, alpha-macroglobulin TED domain, alpha-2-macroglobulin, VIT domain, kringle superfamily, terpenoid cyclases/protein prenyltransferase alpha-alpha toroid, zinc finger, FYVE related, FYVE zinc finger, immunoglobulin-like fold, lipid transport protein, beta-sheet shell, lipovitellin-phosvitin complex, superhelical domain, lipid transport protein, netrin domain, netrin module non-TIMP type, vWf type A and vWf type D domain, the tissue inhibitor of metalloproteinases-like OB-fold, histone H2A/H2B/H3, histone-fold, peptidase S1, PA clam, fibrinogen alpha/beta/gamma chain C-terminal globular domain, fibrinogen-like C-terminal, serine proteases, trypsin family serine active site and histidine active site, sushi/SCR/CCP superfamily, peptidase S1A chymotrypsin family.

Proteomic analysis using LC-MS/MS, identified a range of innate and adaptive immune proteins to be exported in serum-EVs, including in deiminated form, as listed above. This also included a range of complement components, whereof C3 and C5 were detected as deiminated in serum EVs, while in total EV cargo, C1, C3, C4, C5, C6, C7, C8 and C9 were also identified as hits, as well as factor B and factor H. This correlates with previous findings reporting C3 to be deiminated in teleost fish, both in halibut and cod [13,32,33]. Furthermore, a proteomic analysis of deiminated target proteins in halibut serum identified C5, C7, C8 C9 and C1-inhibitor to be deiminated in whole halibut serum [13]. These findings, and the current study, indicate that not all complement components are exported in EVs in deiminated form, and some are found in deiminated form only in whole serum, while being exported in non-deiminated form in serum-EVs. Recent studies assessing protein deimination across the phylogeny tree have indeed identified various complement components as deimination candidates in a range of taxa [14,16,32–35,37,39,40]. Furthermore, C5 has been verified to be a deimination candidate by bacterial

arginine deiminase, allowing for immune modulation of the host and bacterial immune evasion [18].

In the current study we furthermore evaluated by western blotting some key complement proteins identified by LC/MS-MS in EV total protein cargo and deimination-enriched protein cargo. For this purpose, we used halibut-specific antibodies against C3, C4 and pentraxin-like protein, previously developed and described by our group [13,42]. Using western blotting analysis, we verified the presence of C3, C4 and pentraxin-like protein in halibut serum-EVs, showing that these are indeed exported in EVs, as also identified by LC-MS/MS analysis. The C3 antibody also reacted strongly with the F95 enriched protein eluate from the serum EVs, while a lower signal was seen for C4, indicating that C3 is present at higher levels in deiminated form in serum-EVs, compared with C4. Pentraxin-like protein was only observed in total protein cargo of serum-EVs, but not the F95 enriched serum-EV eluate and this corresponds with the LC-MS/MS analysis which revealed hits with a pentraxin for the total protein cargo analysis of serum-EVs, but not the F95-enriched fraction. Our current findings in halibut serum-EV cargo also correspond to our previous analysis on serum EVs and mucus EVs in Atlantic cod, where C3 was detected at higher levels in serum-EVs than C4, both for total protein as well as in the F95-enriched eluate for a putative deiminated form [32,33,38]. Furthermore, cod serum and mucus EVs were also found to contain pentraxin-like protein (CRP-like), which was not detected in deiminated form in the cod EVs, similar to as observed for pentraxin-like protein in halibut serum-EVs in the current study.

Overall, our and others' findings indicate that the complement system can be modulated by deimination both by the host and by pathogen interactions. Understanding of post-translational regulation of complement components via deimination is still in its infancy and requires in depth investigation as deimination may facilitate multifaceted functions of complement proteins in immunity and tissue remodelling in health and disease, also across phylogeny. Such regulation via deimination may furthermore allow for targeted modulation in relation to a range of pathological processes, including infection and autoimmune diseases, where PADs, EVs and the complement system all play important roles.

Besides differences in EV cargo for complement components, proteins that were only identified in whole protein cargo (and not in the F95 eluate) related to a range of innate and adaptive immune factors as well as metabolic and gene regulatory function. These included Apolipoprotein Bb, Apolipoprotein M, Ig-like domain-containing protein, Ig heavy chain Mem5-like, IGv domain-containing protein, Immunoglobulin light chain, nattectin, SERPIN domain-containing protein, Lysozyme, ceruloplasmin, vitellogenin, apoptosis-stimulating of p53 protein 2 Bcl2-binding protein, plasminogen, keratin, type I cytoskeletal 13-like, EGF-like domain-containing protein, hephaestin-like protein 1, desmoglein-2, carboxypeptidase Q, beta 1-globin, antithrombin-III, collagen alpha-1(XII) chain, desmoplakin, biotinidase, collagenase 3, cathepsin L1-like, prothrombin, putative insulin-like growth factor binding protein, sushi domain-containing protein 2 isoform 2, 14_3_3 domain-containing protein, catechol O-methyltransferase domain-containing protein 1, pleckstrin, hyaluronan-binding protein 2, retrotransposon-derived, FH2 domain-containing protein 1-like, protein-tyrosine-phosphatase, thyroid hormone receptor interactor 11, roundabout-like axon guidance receptor protein 2, protein Z-dependent protease inhibitor-like, myosin phosphatase Rho interacting protein, multidrug and toxin extrusion protein, FH2 domain containing 4, nesprin-2, histone H3-trimethyl-L-lysine(9) demethylase, centrosomal protein of 290 kDa and titin-like protein. These all relate to the protein-interaction networks identified for whole EV protein cargo, listed above and shown in Figure 5.

Furthermore, some protein candidates were only detected in the F95 eluate, indicating that they were exported in deiminated form only in the serum EVs. This included cytoplasmic 2 actin, tubulin alpha chain, keratin 93, trypsin-3-like, centrosomal protein of 162 kDa, 2-phospho-D-glycerate hydro-lyase, integrin beta and myosin_tail_1 do-

main-containing protein. Compared with a previous analysis from our group on deiminated proteins in whole halibut serum [13], deiminated candidates found here to be exported specifically in EVs are cytoplasmic 2 actin, tubulin alpha chain, centrosomal protein of 162 kDa, 2-phospho-D-glycerate hydro-lyase, integrin beta and myosin_tail_1 domain-containing protein. This indicates that there are differences in deiminated protein cargo in serum-EVs compared with whole serum, and this corresponds to findings from other comparative studies analysing differences in KEGG (Kyoto encyclopedia of genes and genomes) and GO (gene ontology) enrichment pathways for deiminated proteins in whole serum/plasma versus EVs in diverse taxa, including in cow, camelid, alligator, rat, naked mole-rat, shark and cod [14,16,33–36,43]. Furthermore, differences in deimination signatures in whole serum versus EV cargo have been reported to relate to immune/growth trade off in response to environmental temperature in teleost cod [33]. Such findings, including the findings reported in our current study, emphasise that variations in EV cargo, including via the transport of deiminated proteins, may play hitherto under-recognized and important roles in cellular communication in health and disease across the phylogeny tree.

4. Materials and Methods

4.1. Fish and Sampling

Blood was collected from four adult halibut (*Hippoglossus hippoglossus* L.; weight 4.5–5.0 kg), which were obtained from the experimental fish farm Fiskeldi Eyjafjardar hf, Thorlakshofn, Iceland (under licence from the Institute for Experimental Pathology, University of Iceland, number #0002 kt-650269–4549, approved by the central animal ethics committee in Iceland (Icelandic Food Regulation Authority, MAST Matvælastofnun). Following 1–3 mL blood collection from a gill vessel, the blot was left to clot overnight at 4 °C, and thereafter serum collection was performed by centrifugation at 750 g for 10 min. Serum aliquots of 200 µL were stored at –20 °C until used. The health status of the fish at the fish farm was routinely examined at regular 3 monthly intervals by the Fish Disease Laboratory, Institute for Experimental Pathology, Keldur, Iceland, declaring the fish healthy and disease free.

4.2. EV Isolation and Nanoparticle Tracking (NTA) Analysis

EVs were isolated from halibut serum of four individual fish by step-wise centrifugation, according to previously established methods in our laboratory [14,22,39] and according to the guidelines of the International Society for Extracellular Vesicles (ISEV) [49]. Total EV isolates were prepared from the individual 100 µL serum aliquots ($n = 4$), which were diluted 1:5 in Dulbecco's PBS (DPBS, which had previously been ultrafiltered using a 0.22 µm filter, before use) and then centrifuged at 4000 g for 30 min at 4 °C, to ensure the removal of aggregates and apoptotic bodies. The supernatants containing the EVs were collected and ultracentrifuged at 100,000 g for 1 h at 4 °C. The EV-enriched pellets were then resuspended in 1 mL DPBS ("washing step") and ultracentrifuged again at 100,000 g for 1 h at 4 °C. The final EV-enriched pellets were then resuspended in 100 µL DPBS and analysed by NTA for size distribution profiles, using the NanoSight NS300 system (Malvern Panalytical Ltd, Malvern, UK), recording five 60 s videos for each sample. The number of particles per frame was kept in-between 40 to 60 and replicate histograms were generated from the videos, using the NanoSight software 3.0 (Malvern), representing mean and confidence intervals of the five recordings for each sample.

4.3. Transmission Electron Microscopy (TEM)

EVs were further characterised by Transmission Electron Microscopy (TEM) as follows: A pool of EVs, isolated from serum of the four individual animals as described above, was used for morphological analysis using TEM according to previously de-

scribed methods [16,23,34]. In brief, 100 mM sodium cacodylate buffer (pH 7.4) was used to resuspend the EVs, which were then placed onto a glow discharged carbon support film on a grid and fixed at room temperature for 1 min in 2.5% glutaraldehyde in 100 mM sodium cacodylate buffer (pH 7.0). For the staining of EVs, 2% aqueous Uranyl Acetate (Sigma, Gillingham, UK) was used for 1 min, thereafter removing the excess stain. EV imaging was performed using a JEOL JEM 1400 transmission electron microscope (JEOL, Tokyo, Japan) operated at 80 kV at a magnification of 30,000× to 60,000×. Digital images were recorded using an AMT XR60 CCD camera (Deben, Bury St. Edmunds, UK).

4.4. Proteomic Analysis and Protein Identification

The isolation of deiminated/citrullinated proteins from serum-EVs was carried out by immunoprecipitation, using the Catch and Release® v2.0 Reversible Immunoprecipitation System (Merck, Feltham, UK) according to the manufacturer's instructions in conjunction with the pan-deimination F95 antibody (MABN328, Merck), which specifically detects proteins modified by citrullination [50]. F95 enrichment was performed overnight at 4 °C on a rotating platform from a pool of sera ($n = 4$ individuals), followed by the elution of the F95 bound proteins under reducing conditions, according to the manufacturer's instructions (Merck). The F95 eluate was diluted in 2 × Laemmli sample buffer for subsequent SDS-PAGE and western blotting analysis. The total F95 bound protein eluate, as well as total protein from serum-EVs, were also analysed by liquid chromatography–mass spectrometry (LC-MS/MS) (performed by Cambridge Centre for Proteomics, Cambridge, UK), by in-gel digestion, as previously described [32,34]. For the identification of deiminated protein hits, the files were submitted to the Mascot search algorithm (Matrix Science, London, UK) and searched against the UniProt database for Teleostei (CCP_Teleostei Teleostei_20201009; 4085639 sequences; 2121030378 residues). A search was also conducted against a common contaminant database (cRAP 20190401; 125 sequences; 41,129 residues). A significance threshold value of $p < 0.05$ and a peptide cut-off score of 53 were also applied (carried out by Cambridge Proteomics, Cambridge, UK).

In addition to the LC-MS/MS analysis, both total EV proteins and F95 enriched EV proteins were assessed specifically for halibut C3, C4 and pentraxin-like protein content, using mono-specific antibodies, which were previously prepared against these proteins by our group [13,42] (see Section 4.5).

4.5. Western Blotting

Serum EVs were pooled ($n = 4$), reconstituted 1:1 in 2 × Laemmli sample buffer and boiled at 100 °C for 5 min before separation by SDS-PAGE, using 4–20% TGX gels (Bio-Rad, Watford, UK). Proteins were blotted onto 0.45 µm nitrocellulose membranes (Bio-Rad, UK) using semi-dry transfer for 1 h at 15 V and even protein transfer was assessed using Ponceau S (Sigma, Gillingham, UK) staining. Membranes were blocked for 1 h at RT in 5% bovine serum albumin (BSA, Sigma) in Tris-buffered saline containing Tween20 (TBS-T). Primary antibody incubation was performed overnight at 4 °C on a shaking platform, diluting the antibodies in TBS-T. EVs were assessed for the EV-specific markers Flotillin-1 (ab41927, 1/1000) and CD63 (ab216130, 1/1000). EV cargo was assessed for halibut pentraxin-like protein (1/1000; [13]), halibut C3 (1/1000; [42]) and halibut C4 (1/1000; [13]), using halibut-specific antibodies previously generated in our laboratory [13,42]. Following primary antibody incubation, the membranes were washed three times in TBS-T and then incubated at room temperature for 1 h in either the corresponding anti-mouse IgG (for anti-pentraxin, anti-C3 and anti-C4 antibodies) or anti-rabbit IgG (for CD63 and Flot-1 antibodies) HRP-conjugated secondary antibodies (1/3000; BioRad). Thereafter, the membranes were washed in TBS-T five times for 10 min and then visualised using a UVP BioDoc-IT™ System (Cambridge, UK) in conjunction with ECL (Amersham, Merck, UK).

4.6. Silver Staining

Total proteins isolated from serum EVs and the F95-enriched protein eluates from halibut serum EVs were assessed by silver staining following SDS-PAGE in 4–20% gradient TGX gels (BioRad) under reducing conditions. The BioRad Silver Stain Plus Kit (1610449, BioRad) was used to visualise the protein bands according to the manufacturer's instructions (BioRad).

4.7. Protein–Protein Interaction Network Analysis

For the construction of protein–protein interaction networks for deiminated proteins identified in halibut serum-EVs and for total protein content from serum EVs, respectively, STRING analysis (Search Tool for the Retrieval of Interacting Genes/Proteins; <https://string-db.org/>) was applied. The protein networks were built based on the protein names, using the teleost fish STRING database and using the function of “search multiple proteins”. Settings were as “basic” and “medium confidence”. Colour lines connecting the nodes represent the following evidence-based interactions for the network edges: “known interactions” (this is based on experimentally determined data or curated databases); “predicted interactions” (this is based on gene co-occurrence, gene neighbourhood or gene fusion); “others” (this is based on co-expression, text mining or protein homology (see colour key for lines in Figure 4A). Networks were assessed for local network clusters, reactome pathways, PFAM and SMART protein domains and UniProt keywords. The zebrafish (*Danio rerio*) STRING database was used as representative for Teleostei for the creation of the networks as no specific halibut STRING database is available (due to lack of annotation available for halibut), and *D. rerio* showed the most hit number identity with the proteins identified in halibut EVs.

4.8. Statistical Analysis

The Nanosight 3.0 software (Malvern) was used for the generation of NTA curves, which represent mean and standard error of mean (SEM), indicated by confidence intervals. Significance for protein network analysis generated in STRING (<https://string-db.org/>) was considered as $p \leq 0.05$.

5. Conclusions

This study is the first report of EV profile signatures in halibut, analysing total protein and specifically also deiminated protein cargo in serum-EVs. Halibut serum EVs showed a poly-dispersed population with EVs in the size range of 50–600 nm, positive for phylogenetically conserved EV markers. Proteomic analysis of EV total protein cargo revealed 124 protein hits and 37 deiminated protein hits, whereof 15 hits were particularly identified in deiminated form only. Protein interaction network analysis revealed GO pathways for EV mediated protein cargo transport, relating to a range of gene regulatory, immune, metabolic and developmental processes, some of which were enriched for deiminated proteins. Further assessment of key immune related proteins—complement components C3, C4 and pentraxin—identified that C3 is exported in serum-EVs at higher levels than C4, also in deiminated form, while pentraxin was found in whole protein EV content only, but not in deiminated form. Our findings emphasize the putative differences in cell communication mediated by EV protein versus post-translationally deiminated protein cargo (the “EV-citrullinome”), providing novel insights into EV-mediated communication in halibut serum. Our findings furthermore contribute to current understanding of EV signatures across the phylogeny tree, with the potential for biomarker development and EV “fingerprinting” for the assessment of animal health.

Supplementary Materials: The following are available online at www.mdpi.com/1422-0067/22/2/875/s1, Table S1: Deiminated proteins in serum-EVs of halibut (*Hippoglossus hippoglossus* L), as

identified by F95-enrichment in conjunction with LC-MS/MS analysis; full LC-MS/MS data. Table S2: Total proteins in serum-EVs of halibut (*Hippoglossus hippoglossus* L.); full LC-MS/MS data.

Author Contributions: Conceptualization, S.L.; methodology, S.L. A.W.D. and I.K.; validation, B.M., I.K., A.W.D. and S.L.; formal analysis, I.K. and S.L.; investigation, B.M. and S.L.; resources, B.M., I.K., A.W.D. and S.L.; data curation, S.L.; writing—original draft preparation, S.L.; writing—review and editing, B.M., A.W.D. and S.L.; visualization, I.K. and S.L.; supervision, S.L.; project administration, S.L.; funding acquisition, S.L. All authors have read and agreed to the published version of the manuscript.

Funding: This work is partly based on previous support by the EC grant Fishaid QLK2-CT-2000-01076 (B.M. and S.L.), the Icelandic Ministry of Fisheries (to S.L.), the Icelandic Research Council (to S.L.) and the European Molecular Biology Organization (to S.L.). The work was furthermore supported by internal funding from the University of Westminster (to S.L.).

Institutional Review Board Statement: The study was conducted under license from the Institute for Experimental Pathology, University of Iceland, number #0002 kt-650269–4549, approved by the central animal ethics committee in Iceland (Icelandic Food Regulation Authority, MAST Matvælastofnun).

Data Availability Statement: Data is contained within the article and supplementary material.

Acknowledgments: The authors wish to thank Birgir Kristjánsson and the staff at Fiskeldi Eyjafjardar, Þorlákshöfn, Iceland, and the staff at Fiskey hf, Hjalteyri, Iceland for providing the halibut and sampling facilities. Thanks to Michael Deery and Yagnesh Umrana at the Cambridge Centre for Proteomics for performing the LC-MS/MS analysis. Thanks are due to The Guy Foundation for funding the purchase of equipment utilised in this work.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Russel, F.S. *The Eggs and Planktonic Stages of British Marine Fishes*; Academic Press: London, UK, 1976; ISBN 0-12-604050-8.
2. Mangor-Jensen, A.; Harboe, T.; Shields, R.J.; Gara, B.; Naas, K.E. Atlantic halibut, *Hippoglossus hippoglossus* L.; larvae cultivation literature, including a bibliography. *Aquac. Res.* **1998**, *29*, 857–886.
3. Vossenaar, E.R.; Zendman, A.J.; van Venrooij, W.J.; Puijn, G.J. PAD, a growing family of citrullinating enzymes: Genes, features and involvement in disease. *Bioessays* **2003**, *25*, 1106–1118.
4. Wang, S.; Wang, Y. Peptidylarginine deiminases in citrullination, gene regulation, health and pathogenesis. *Biochim. Biophys. Acta* **2013**, *1829*, 1126–1135.
5. Witalison, E.E.; Thompson, P.R.; Hofseth, L.J. Protein Arginine Deiminases and Associated Citrullination: Physiological Functions and Diseases Associated with Dysregulation. *Curr. Drug Targets* **2015**, *16*, 700–710.
6. György, B.; Toth, E.; Tarcsa, E.; Falus, A.; Buzas, E.I. Citrullination: A posttranslational modification in health and disease. *Int. J. Biochem. Cell Biol.* **2006**, *38*, 1662–1677.
7. Bicker, K.L.; Thompson, P.R. The protein arginine deiminases: Structure, function, inhibition, and disease. *Biopolymers* **2013**, *99*, 155–163.
8. Tarcsa, E.; Marekov, L.N.; Mei, G.; Melino, G.; Lee, S.C.; Steinert, P.M. Protein unfolding by peptidylarginine deiminase. Substrate specificity and structural relationships of the natural substrates trichohyalin and filaggrin. *J. Biol. Chem.* **1996**, *271*, 30709–30716.
9. Jeffrey, C.J. Protein moonlighting: What is it, and why is it important? *Philos. Trans. R. Soc. B Biol. Sci.* **2018**, *373*, 20160523.
10. Nomura, K. Specificity and mode of action of the muscle-type protein-arginine deiminase. *Arch. Biochem. Biophys.* **1992**, *293*, 362–369.
11. Rebl, A.; Köllner, B.; Anders, E.; Wimmers, K.; Goldammer, T. Peptidylarginine deiminase gene is differentially expressed in freshwater and brackish water rainbow trout. *Mol. Biol. Rep.* **2010**, *37*, 2333–2339.
12. Magnadóttir, B.; Hayes, P.; Hristova, M.; Bragason, B.P.; Nicholas, A.P.; Dodds, A.W.; Guðmundsdóttir, S.; Lange, S. Post-translational Protein Deimination in Cod (*Gadus morhua* L.) Ontogeny—Novel Roles in Tissue Remodelling and Mucosal Immune Defences? *Dev. Comp. Immunol.* **2018**, *87*, 157–170.
13. Magnadóttir, B.; Bragason, B.T.; Bricknell, I.R.; Bowden, T.; Nicholas, A.P.; Hristova, M.; Guðmundsdóttir, S.; Dodds, A.W.; Lange, S. Peptidylarginine deiminase and deiminated proteins are detected throughout early halibut ontogeny—Complement components C3 and C4 are post-translationally deiminated in halibut (*Hippoglossus hippoglossus* L.). *Dev. Comp. Immunol.* **2019**, *92*, 1–19.

14. Criscitiello, M.F.; Kraev, I.; Lange, S. Deiminated proteins in extracellular vesicles and plasma of nurse shark (*Ginglymostoma cirratum*)—Novel insights into shark immunity. *Fish Shellfish Immunol.* **2019**, *92*, 249–255.
15. Lange, S.; Gögel, S.; Leung, K.Y.; Vernay, B.; Nicholas, A.P.; Causey, C.P.; Thompson, P.R.; Greene, N.D.; Ferretti, P. Protein deiminases: New players in the developmentally regulated loss of neural regenerative ability. *Dev. Biol.* **2011**, *355*, 205–214.
16. Criscitiello, M.F.; Kraev, I.; Petersen, L.H.; Lange, S. Deimination Protein Profiles in *Alligator mississippiensis* Reveal Plasma and Extracellular Vesicle-Specific Signatures Relating to Immunity, Metabolic Function, and Gene Regulation. *Front. Immunol.* **2020**, *11*, 651.
17. Novák, L.; Zubáčová, Z.; Karnkowska, A.; Kolisko, M.; Hroudová, M.; Stairs, C.W.; Simpson, A.G.; Keeling, P.J.; Roger, A.J.; Čepička, I.; et al. Arginine deiminase pathway enzymes: Evolutionary history in metamonads and other eukaryotes. *BMC Evol. Biol.* **2016**, *16*, 197.
18. Bielecka, E.; Scavenius, C.; Kantyka, T.; Jusko, M.; Mizgalska, D.; Szmigielski, B.; Potempa, B.; Enghild, J.J.; Prossnitz, E.R.; Blom, A.M.; et al. Peptidyl arginine deiminase from *Porphyromonas gingivalis* abolishes anaphylatoxin C5a activity. *J. Biol. Chem.* **2014**, *289*, 32481–32487.
19. Kosgodage, U.S.; Matewale, P.; Mastroianni, G.; Kraev, I.; Brotherton, D.; Awamaria, B.; Nicholas, A.P.; Lange, S.; Inal, J.M. Peptidylarginine deiminase inhibitors reduce bacterial membrane vesicle release and sensitize bacteria to antibiotic treatment. *Front. Cell. Infect. Microbiol.* **2019**, *9*, 227.
20. El-Sayed, A.S.A.; Shindia, A.A.; Abouzaid, A.A.; Yassin, A.M.; Ali, G.S.; Sitohy, M.Z. Biochemical characterization of peptidylarginine deiminase-like orthologs from thermotolerant *Emericella dentata* and *Aspergillus nidulans*. *Enzyme Microb. Technol.* **2019**, *124*, 41–53.
21. Gavinho, B.; Sabatke, B.; Feijoli, V.; Rossi, I.V.; da Silva, J.M.; Evans-Osses, I.; Palmisano, G.; Lange, S.; Ramirez, M.I. Peptidylarginine deiminase inhibition abolishes the production of large extracellular vesicles from *Giardia intestinalis*, affecting host-pathogen interactions by hindering adhesion to host cells. *Front. Cell. Infect. Microbiol.* **2020**, *10*, 417.
22. Bowden, T.J.; Kraev, I.; Lange, S. Extracellular vesicles and post-translational protein deimination signatures in haemolymph of the American lobster (*Homarus americanus*). *Fish Shellfish Immunol.* **2020**, *106*, 79–102.
23. Bowden, T.J.; Kraev, I.; Lange, S. Post-translational protein deimination signatures and extracellular vesicles (EVs) in the Atlantic horseshoe crab (*Limulus polyphemus*). *Dev. Comp. Immunol.* **2020**, *110*, 103714.
24. Bowden, T.J.; Kraev, I.; Lange, S. Extracellular Vesicles and Post-Translational Protein Deimination Signatures in Mollusca—The Blue Mussel (*Mytilus edulis*), Soft Shell Clam (*Mya arenaria*), Eastern Oyster (*Crassostrea virginica*) and Atlantic Jackknife Clam (*Ensis leei*). *Biology* **2020**, *9*, 416.
25. Inal, J.M.; Ansa-Addo, E.A.; Lange, S. Interplay of host-pathogen microvesicles and their role in infectious disease. *Biochem. Soc. Trans.* **2013**, *41*, 258–262.
26. Colombo, M.; Raposo, G.; Théry, C. Biogenesis, secretion, and intercellular interactions of exosomes and other extracellular vesicles. *Annu. Rev. Cell Dev. Biol.* **2014**, *30*, 255–289.
27. Kholia, S.; Jorfi, S.; Thompson, P.R.; Causey, C.P.; Nicholas, A.P.; Inal, J.M.; Lange, S. A novel role for peptidylarginine deiminases in microvesicle release reveals therapeutic potential of PAD inhibition in sensitizing prostate cancer cells to chemotherapy. *J. Extracell. Vesicles* **2015**, *4*, 26192.
28. Kosgodage, U.S.; Uysal-Onganer, P.; MacLachy, A.; Kraev, I.; Chatterton, N.P.; Nicholas, A.P.; Inal, J.M.; Lange, S. Peptidylarginine Deiminases Post-Translationally Deiminate Prohibitin and Modulate Extracellular Vesicle Release and MicroRNAs in Glioblastoma Multiforme. *Int. J. Mol. Sci.* **2018**, *20*, 103.
29. Uysal-Onganer, P.; MacLachy, A.; Mahmoud, R.; Kraev, I.; Thompson, P.R.; Inal, J.M.; Lange, S. Peptidylarginine Deiminase Isozyme-Specific PAD2, PAD3 and PAD4 Inhibitors Differentially Modulate Extracellular Vesicle Signatures and Cell Invasion in Two Glioblastoma Multiforme Cell Lines. *Int. J. Mol. Sci.* **2020**, *21*, 1495.
30. Turchinovich, A.; Drapkina, O.; Tonevitsky, A.; Transcriptome of extracellular vesicles: State-of-the-art. *Front. Immunol.* **2018**, *10*, 202.
31. Vagner, T.; Chin, A.; Mariscal, J.; Bannykh, S.; Engman, D.M.; di Vizio, D. Protein composition reflects extracellular vesicle heterogeneity. *Proteomics* **2019**, *19*, e1800167.
32. Magnadóttir, B.; Kraev, I.; Guðmundsdóttir, S.; Dodds, A.W.; Lange, S. Extracellular vesicles from cod (*Gadus morhua* L.) mucus contain innate immune factors and deiminated protein cargo. *Dev. Comp. Immunol.* **2019**, *99*, 103397.
33. Magnadóttir, B.; Uysal-Onganer, P.; Kraev, I.; Dodds, A.W.; Guðmundsdóttir, S.; Lange, S. Extracellular vesicles, deiminated protein cargo and microRNAs are novel serum biomarkers for environmental rearing temperature in Atlantic cod (*Gadus morhua* L.). *Aquac. Rep.* **2020**, *16*, 100245.
34. Criscitiello, M.F.; Kraev, I.; Lange, S. Deiminated proteins in extracellular vesicles and serum of llama (*Lama glama*)—Novel insights into camelid immunity. *Mol. Immunol.* **2020**, *117*, 37–53.
35. Criscitiello, M.F.; Kraev, I.; Lange, S. Post-translational protein deimination signatures in serum and serum-extracellular vesicles of *Bos taurus* reveal immune, anti-pathogenic, anti-viral, metabolic and cancer-related pathways for deimination. *Int. J. Mol. Sci.* **2020**, *21*, 2861.
36. Pamenter, M.E.; Uysal-Onganer, P.; Huynh, K.W.; Kraev, I.; Lange, S. Post-translational deimination of immunological and metabolic protein markers in plasma and extracellular vesicles of naked mole-rat (*Heterocephalus glaber*). *Int. J. Mol. Sci.* **2019**, *20*, 5378.
37. Phillips, R.A.; Kraev, I.; Lange, S. Protein deimination and extracellular vesicle profiles in Antarctic seabirds. *Biology* **2020**, *9*, 15.

38. Lange, S.; Kraev, I.; Magnadóttir, B.; Dodds, A.W. Complement component C4-like protein in Atlantic cod (*Gadus morhua* L.)—Detection in ontogeny and identification of post-translational deimination in serum and extracellular vesicles. *Dev. Comp. Immunol.* **2019**, *101*, 103437.
39. Magnadóttir, B.; Uysal-Onganer, P.; Kraev, I.; Svansson, V.; Hayes, P.; Lange, S. Deiminated proteins and extracellular vesicles—Novel serum biomarkers in whales and orca. *Comp. Biochem. Physiol. Part D Genom. Proteom.* **2020**, *34*, 100676.
40. Magnadóttir, B.; Uysal-Onganer, P.; Kraev, I.; Svansson, V.; Skirnisson, K.; Lange, S. Deiminated proteins and extracellular vesicles as novel biomarkers in pinnipeds: Grey seal (*Halichoerus grypus*) and harbour seal (*Phoca vitulina*). *Biochimie* **2020**, *171*–172, 79–90.
41. Iliev, D.; Strandskog, G.; Nepal, A.; Aspar, A.; Olsen, R.; Jørgensen, J.; Wolfson, D.; Ahluwalia, B.S.; Handzhiyski, J.; Mironova, R. Stimulation of exosome release by extracellular DNA is conserved across multiple cell types. *FEBS J.* **2018**, *285*, 3114–3133.
42. Lange, S.; Dodds, A.W.; Magnadóttir, B. Isolation and characterization of complement component C3 from Atlantic cod (*Gadus morhua* L.) and Atlantic halibut (*Hippoglossus hippoglossus* L.). *Fish Shellfish Immunol.* **2004**, *16*, 227–239.
43. Sancandi, M.; Uysal-Onganer, P.; Kraev, I.; Mercer, A.; Lange, S. Protein Deimination Signatures in Plasma and Plasma-EVs and Protein Deimination in the Brain Vasculature in a Rat Model of Pre-Motor Parkinson's Disease. *Int. J. Mol. Sci.* **2020**, *21*, 2743.
44. Kosgodage, U.S.; Matewele, P.; Awamaria, B.; Kraev, I.; Warde, P.; Mastroianni, G.; Nunn, A.V.; Guy, G.W.; Bell, J.D.; Inal, J.M.; et al. Cannabidiol Is a Novel Modulator of Bacterial Membrane Vesicles. *Front. Cell. Infect. Microbiol.* **2019**, *9*, 324.
45. Zhang, X.; Hubal, M.J.; Kraus, V.B. Immune cell extracellular vesicles and their mitochondrial content decline with ageing. *Immun. Ageing* **2020**, *17*, 1.
46. Faught, E.; Henrickson, L.; Vijayan, M.M. Plasma exosomes are enriched in Hsp70 and modulated by stress and cortisol in rainbow trout. *J. Endocrinol.* **2017**, *232*, 237–246.
47. Cadonic, I.G.; Ikert, H.; Craig, P.M. Acute air exposure modulates the microRNA abundance in stress responsive tissues and circulating extracellular vesicles in rainbow trout (*Oncorhynchus mykiss*). *Comp. Biochem. Physiol. Part D Genom. Proteom.* **2020**, *34*, 100661.
48. Antwi-Baffour, S.; Malibha-Pinchbeck, M.; Stratton, D.; Jorfi, S.; Lange, S.; Inal, J. Plasma mEV levels in Ghanain malaria patients with low parasitaemia are higher than those of healthy controls, raising the potential for parasite markers in mEVs as diagnostic targets. *J. Extracell. Vesicles* **2019**, *9*, 1697124.
49. Théry, C.; Witwer, K.W.; Aikawa, E.; Alcaraz, M.J.; Anderson, J.D.; Andriantsitohaina, R.; Antoniou, A.; Arab, T.; Archer, F.; Atkin-Smith, G.K.; et al. Minimal information for studies of extracellular vesicles 2018 (MISEV2018): A position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *J. Extracell. Vesicles* **2018**, *7*, 1535750.
50. Nicholas, A.P.; Whitaker, J.N. Preparation of a monoclonal antibody to citrullinated epitopes: Its characterization and some applications to immunohistochemistry in human brain. *Glia* **2002**, *37*, 328–336.